Today

- q-ary codes.
- Algebraic-geometry & Codes.
- Proof of concept.
- Statement of big claim.

Hamming balls in $\mathbb{F}_q^n$

- Let $H_q(p) = p \log_q \frac{q-1}{p} + (1-p) \log_q \frac{1}{1-p}$.
- Let $\mathrm{Vol}_q(r,n) =$ Volume of Hamming ball of radius r in $\mathbb{F}_q^n$.
- Then $\mathrm{Vol}_q(r,n) = q^{(H_q(p)+o(1))n}$.
- q-ary GV bound:
  Theorem: There exists an infinite family of q-ary codes of rate R and relative distance δ satisfying

$$R \geq 1 - H_q(\delta)$$

.

Understanding q-ary GV bound

- To get a sense of $H_q(\delta)$, fix $0 < \delta < 1$ and let $q \to \infty$. Get

$$R \geq 1 - \delta - H_2(\delta)/\log q - o(1/\log q)$$

- Contrast with Singleton (Project on to first $k-1$ coordinates) upper bound on rate:

$$R \leq 1 - \delta$$

- I.e., GV bound approaches Singleton bound at logarithmic rate in q.
- Is this best possible? RS codes achieve Singleton bound and q is pretty small!

- Today: Algebraic-geometry (AG) codes: Achieve

$$R \geq 1 - \delta - \frac{1}{\sqrt{q} - 1}$$

  - Needs q square and prime power.
  - Clearly better for large q.
  - In fact, better for $q \geq 49$.

Algebraic-geometry codes

- Conceived by Goppa in late 70's - early 80's.
- 1982 - Surprising breakthrough ...
  - Due to Tsfasman, Vladuts, Zink.
  - Based on some prior work of Ihara.
  - Codes better than random for suff. large, but constant sized, alphabet.
- Almost unique in history of explicit constructions ....

Motivation: Bivariate Codes

- Consider codes obtained by evaluations of bivariate polynomials $Q(x, y)$ of deg. $\leq l$ in each variable.
- Gives $\left[q^2, l^2, \left(1 - \frac{l}{q}\right)^2\right]_q$ code.
- Contrast w. $\left[q^2, l^2, q^2 - l^2\right]_{q^2}$ RS code.
  - Bivariate alphabet smaller.
  - Distance smaller by $2l(q - l)$.
- Why this $q - l$ deficit?
  - On axis-parallel line l points zero imply q points zero.
  - For every line defect of $q - l$.

AG code idea

- Don't evaluate poly on all points on plane.
- Ideally, don't use more than l points on line.
- Pragmatically, don't use much more than l points on line.
- But there exist other bad examples. Degree 2 curves, Degree 3 curves.
- So, don't use too many points on any (low-degree) curve.
- How to find such points? Use points on some low-degree curve.

Algebraic curves in the plane

Defn: Given a bivariate polynomial $R(x, y)$ of total degree D, the set of points

$$\{(a, b) \in \Sigma^2 \mid R(a, b) = 0\}$$

is called an algebraic curve of degree D in the plane.

Basic result from algebraic geometry:
Nice algebraic curves don't meet other nice algebraic curves very often.

Bezout's Thm: Curves R_1, R_2 of deg. D_1, D_2 share at most $D_1 D_2$ common zeroes.

Example (stolen from Shokrollahi)

- Let $q = 13$
  $R(x, y) = y^2 - 2(x-1)x(x+1)$.
- Code obtained by evaluating (certain) polynomials at zeroes of R.
- Fact: There exist 19 zeroes of R.
- Legal polynomials: linear combinations of $\{1, x, y, x^2, xy, x^3\}$.
- If legal poly has 6 zeroes, then it is identically zero.
- Gives $[19, 6, 13]_{13}$ code.
  (RS would give $[19, 6, 14]_{19}$ code.)

Codes from Planar Curves

- Generally:
  - Evaluating polys of deg. $\leq l$
  - At zeroes of R, irreducible, of degree D, with n zeroes.
  - Gives $[n, k, n - Dl]_q$ code,
    for $k = \begin{cases} \binom{l+2}{2} & \text{if } l < D \\ \binom{l+2}{2} - \binom{l-D+2}{2} & \text{if } l \geq D \end{cases}$.
- Distance by Bezout's theorem.

Finding good curves

How to find R with large n?

- No general method.
- But some well-known curves do well. e.g. Hermitian curve for $q = r^2$:
  - $x^{r+1} - y^r - y = 0$
  - has $r^3 + 1$ points.
  - Gives $[r^3+1, \binom{r+2}{2}, r^3+1-(r)(r+1)]_{r^2}$ code.
- Bivariate polys gave
  $[r^4, \binom{r+2}{2}, r^4 - r^3]_{r^2}$.

Going to Higher Dimension

- So far, went from alphabet n to (at best) $\sqrt{n}$.
- To do better need more variables.
- General AG codes:
  - Pick m variables.
  - Put $m - 1$ polynomial constraints.
  - Evaluate polynomials on zeroes.

"State-of-the-art" codes

[Garcia & Stichtenoth]

- $q = r^2$.
- Variables $x_1, \ldots, x_m, y_1, \ldots, y_m$.
- Constraints:
  $x_1^{r+1} = y_1^r + y_1$.
  $x_2 x_1 = y_1$.
  $x_2^{r+1} = y_2^r + y_2$.
  $\vdots$
  $x_m x_{m-1} = y_{m-1}$.
  $x_m^{r+1} = y_m^r + y_m$.
- # zeroes $\geq (r^2 - 1) r^m$.

Keeping track of distance

- Bezout's theorem becomes weak.
- Polynomials ordered by "order".
  Order axioms:
  - $\mathrm{ord}(f+g) \leq \max\{\mathrm{ord}(f), \mathrm{ord}(g)\}$.
  - $\mathrm{ord}(f * g) = \mathrm{ord}(f) + \mathrm{ord}(g)$.
  - f has at most $\mathrm{ord}(f)$ zeroes.
  - Polynomials of all except g orders exist.
  - g = genus of curve.
  - Genus of Garcia-Stichtenoth curve $\leq (r+1)r^m$.
- AG codes follow.

Summary: RS vs. AG

	RS	AG
Coordinates	$\mathbb{F}_q$	Points on curves
Messages	Polynomials deg $< k$	Polynomials order $< k$
Encoding	Evaluations	Evaluations
Distance	$n-k+1$	$n-k+1$
Dimension	k	$k-$ genus
Axioms	zeroes $\leq$ deg. Sum rule Product rule dim. $>$ deg.	zeroes $\leq$ order Sum rule Product rule dim. $>$ order $-$ g

Computational requirements

- Classical AG codes computable in $O(n^{30})$ time.
- Newer AG codes computable in $O(n^{17})$ time.
- Rumors of $O(n^2)$ time computability.
- Belief in explicit constructions.

Some best known codes

Fix $q = 2$. Given k and $d/n = \frac{1}{2} - \epsilon$, what is the best known code? (Will allow $\epsilon = \epsilon(n)$).

- Random code: $n = O(\frac{k}{\epsilon^2})$.
- RS $\circ$ Hadamard: $n = \frac{k^2}{\epsilon^2}$.
- AG $\circ$ Hadamard: $n = O(\frac{k}{\epsilon^3 \log(1/\epsilon)})$.
- [ABNNR]: $n = O(\frac{k}{\epsilon^3})$. (Polylog space constructible).