

- **Reading:** Gallian Chapter 5

## 1 Permutation Groups: Basics

- **Def:** A *permutation group* on a set  $A$  is a subgroup of  $Sym(A)$  (the set of permutations of  $A$  under composition).

- **Examples:**

- $S_n$
- $D_n$  (two choices for  $A$ )
- $GL_n(\mathbb{R})$

[Technically,  $D_n$  and  $GL_n(\mathbb{R})$  are only “isomorphic” to permutation groups on  $[n]$  and  $\mathbb{R}^n$ , respectively.]

- Today we’ll focus on  $A = [n] = \{1, \dots, n\}$ , ie  $S_n$  and its subgroups.
- Running examples:  $\sigma, \tau \in S_7$  defined by

$$\sigma(1) = 5, \sigma(2) = 3, \sigma(3) = 6, \sigma(4) = 7, \sigma(5) = 1, \sigma(6) = 2, \sigma(7) = 4,$$

and

$$\tau(1) = 1, \tau(2) = 2, \tau(3) = 3, \tau(4) = 6, \tau(5) = 7, \tau(6) = 5, \tau(7) = 4.$$

- **Array notation:**

$$\begin{aligned}\sigma &= \begin{bmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ 5 & 3 & 6 & 7 & 1 & 2 & 4 \end{bmatrix} \\ \tau &= \begin{bmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ 1 & 2 & 3 & 6 & 7 & 5 & 4 \end{bmatrix} \\ \tau \circ \sigma &= \begin{bmatrix} 1 & 2 & 3 & 4 & 5 & 6 & 7 \\ 1 & 2 & 3 & 6 & 7 & 5 & 4 \end{bmatrix}\end{aligned}$$

## 2 Cycle Notation

- **Def:** An  $m$ -cycle is a permutation  $\alpha$  for which there exist distinct  $i_1, \dots, i_m$  such that  $\alpha(i_1) = i_2, \alpha(i_2) = i_3, \dots, \alpha(i_{m-1}) = i_m, \alpha(i_m) = i_1$ , and  $\alpha(j) = j$  for all  $j \notin \{i_1, \dots, i_m\}$ .
- **Cycle notation:**  $\alpha = (i_1 i_2 \dots i_m) = (i_2 i_3 \dots i_m i_1) = \dots$

- **Examples:**
- **Q:** What is the order of an  $m$ -cycle?
- **Thm 5.1+:** Every permutation in  $S_n$  can be written as a product of one or more *disjoint* cycles, whose union includes all elements of  $[n]$ . This representation is unique up to the order of the cycles (and cyclic shifts when writing the cycles).
  - We usually don't write the 1-cycles!
- **Proof by example:**  $\sigma =$
- **Graphical view:** View a permutation as a directed graph in which every vertex has indegree and outdegree 1 (possibly with self-loops). Such a graph consists of disjoint cycles.
- **Q (Thm 5.3):** How can we calculate the order of a permutation in terms of its cycles?
- **Example:**  $\text{order}(\sigma) =$
- **Proof in general:**

### 3 Transpositions

- **Def:** A *transposition* is a 2-cycle.
- **Thm 5.4:** Every permutation can be written as a product of transpositions.
  - Not uniquely!
- **Proof:**
- **Thm 5.5+:**
  1. (Even permutations) If a permutation  $\sigma$  has an even number of even-length cycles in disjoint cycle notation, then  $\sigma$  can only be written as product of an even number of transpositions. In such a case,  $\sigma$  is called an *even permutation*.

2. (Odd permutations) If a permutation  $\sigma$  has an odd number of even-length cycles in disjoint cycle notation, then  $\sigma$  can only be written as product of an odd number of transpositions. In such a case,  $\sigma$  is called an *odd permutation*.
- **Proof:** (different from book) Show by induction on  $n$  that if  $\sigma = \alpha_1 \cdots \alpha_n$  for transpositions  $\alpha_i$ , then the parity of the number of even-length cycles in  $\sigma$  equals the parity of  $n$ .
    - Base case ( $n = 0$ ):  $\sigma$  consists of zero even-length cycles.
    - Induction step: Consider what happens when we multiply a permutation  $\sigma = \alpha_1 \cdots \alpha_n$  by an additional transposition  $\alpha_{n+1}$ . Let's do a case analysis depending on how  $\alpha_{n+1} = (ij)$  intersects the disjoint cycles of  $\sigma$ .
      - \* Case 1:  $i$  and  $j$  are both within the same cycle.
      - \* Case 2:  $i$  and  $j$  are within different cycles.
  - **Cor:** The set of even permutations in  $S_n$  is a subgroup, called the *alternating group*  $A_n$ .
  - **Q:** What is  $|A_n|$ ?