

Problem Set 0

Assigned: Tue. Sept. 4, 2018

Due: Fri. Sept. 14, 2018 (5:00 sharp)

- This problem set is optional and will not count for your grade. However, if you have not taken a prior proof-based math course, it is strongly encouraged that you complete and turn in the problem set for practice and feedback on doing proofs.
- You must submit your problem sets electronically on course Canvas site. If you use L^AT_EX, please submit both the source (.tex) and the compiled file (.pdf). Name your files PS0-yourlastname.

Problem 1. (Proof by Contradiction) Prove that $\log_2 6$ is an irrational number.

Problem 2. (Set Equality) Which of the following is true? Prove your answers.

- For every three sets A, B, C , we have $A \cup (B \cap C) = (A \cup B) \cap C$.
- For every three sets A, B, C , we have $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$.

Problem 3. (Induction) The Fibonacci numbers F_0, F_1, \dots are defined inductively by $F_0 = 0$, $F_1 = 1$, and $F_{n+2} = F_{n+1} + F_n$ for all $n \geq 1$. Thus the sequence (starting at F_0) is 0, 1, 1, 2, 3, 5, 8, 13, 21, \dots . Prove by induction that for $n \geq 2$, $F_n \geq \varphi^{n-2}$, where $\varphi = (1 + \sqrt{5})/2$ is the golden ratio.

Problem 4. (Incorrect Induction) What is the wrong with the following proof by induction?

Claim: In every set of n students, all students have the same height.

“Proof” by Induction:

- Base Case: For every set of size 1, the claim is clearly true (all the students in that set have the same height).
- Induction Step: Assume that the claim is true for sets of k students (this is the induction hypothesis), and we’ll prove that it also holds for sets of $k + 1$ students. Consider an arbitrary set S consisting of $k + 1$ students, say $S = \{p_1, \dots, p_{k+1}\}$. Let $S' = \{p_1, \dots, p_k\}$. Since $|S'| = k$, our induction hypothesis tells us that all students in S' have the same height. So now we only need to show that p_{k+1} has the same height too. To do this, consider the set $S'' = \{p_2, \dots, p_{k+1}\}$. Since $|S''| = k$, the induction hypothesis also tells us that all students in S'' have the same height. In particular, p_{k+1} has the same height as p_2 , and hence the same height as all students in S' .