

Marginal Mean Models for Dynamic Regimes

S. A. MURPHY, M. J. VAN DER LAAN, J. M. ROBINS AND CPPRG¹

A dynamic treatment regime is a list of rules for how the level of treatment will be tailored through time to an individual's changing severity. In general, individuals who receive the highest level of treatment are the individuals with the greatest severity and need for treatment. Thus there is planned selection of the treatment dose. In addition to the planned selection mandated by the treatment rules, the use of staff judgment results in unplanned selection of the treatment level. Given observational longitudinal data or data in which there is unplanned selection of the treatment level, the methodology proposed here allows the estimation of a mean response to a dynamic treatment regime under the assumption of sequential randomization.

KEY WORDS: dynamic treatment regimes, nondynamic treatment regimes, causal inference, confounding.

1. INTRODUCTION

Dynamic treatment regimes are individually tailored treatments that provide treatment to individuals only when and if they need the treatment and adjust the level of treatment to the individual's need. In a dynamic regime rules for how the treatment level and type should vary with time are specified prior to the beginning of treatment; these rules are based on time varying measurements of subject-specific need for treatment. Thus the assignment to treatment depends on past individual information. Dynamic treatment regimes are potentially attractive to public policy makers because they treat only subjects who show need for treatment, freeing public and private funds for more intensive treatment of the needy.

The goal of this paper is to provide an estimation method for the mean response that would have been observed had, contrary to fact, the entire study population followed a particular dynamic regime. The available data are observational; that is in the available data, the treatment level may depend on past individual information that is not used in or that is not used according to the particular dynamic regime rules. We employ a structural model, thus the model is a model for the marginal mean of counterfactual or potential responses.

¹ University of Michigan, University of California at Berkeley and Harvard University. CPPRG is the Conduct Problems Prevention Research Group, members include K.L. Bierman (Pennsylvania State Univ.), J.D. Coie (Duke Univ.), M.T. Greenberg (Pennsylvania State Univ.), J.E. Lochman (Univ. of Alabama), R.J. McMahon (Univ. of Washington), and E. Pinderhughes (Vanderbilt Univ.) The research of the first author is partially supported by NIDA grant A P50 DA 10075, DMS 9802885, and SBR 9811983. The research of the second author is partially supported by NIAID grant 1R01AI46182. The research for the third author is partially supported by NIAID grant RO1AI32475.

This work is motivated by the Fast Track prevention program. This is an ongoing randomized trial of a complex preventive intervention versus control. The intervention was designed to prevent the emergence of and reduce the level of conduct disorders and drug use in children at risk due to elevated behavior problems (Bierman et al., 1996; CPPRG, 1999ab; McMahon et al., 1996). Part of the intervention involved implementation of a dynamic treatment regime designed to improve family functioning. At the end of each semester, beginning with the spring semester of first grade, the family counselor filled out a 6 item “home visiting process measure,” describing the quality of parenting and family functioning. Based on the home visiting process measure total score, home visiting assignments for the next semester were made. A simplified version is as follows. A score of 17 or greater corresponds to an assignment of 4 home visits, a score from 9 to 16 corresponds to an assignment of 8 home visits and a score of 8 or below corresponds to an assignment of 16 home visits during the following semester. Thus the rule for assigning dose was

$$d_t(S_{t-1}) = 16I\{S_{t-1} \leq 8\} + 8I\{9 \leq S_{t-1} \leq 16\} + 4I\{17 \leq S_{t-1}\}, t = 1, 2, 3, 4$$

where S_t is an assessment of severity, the home visiting process measure score, at the end of the t th semester, with low values indicating greater severity. Staff were told that in exceptional cases they might need to deviate from the rule for assigning home visiting level; in practice, staff assignments deviated from the rule for approximately 50% of the intervention children. In Section 6, our goal is to estimate what the marginal effect of the above dynamic treatment regime would have been, had staff followed the rules exactly. We demonstrate that by identifying the intervention group (with home visiting assignments as implemented) as an observational study, one can use the methods developed here to achieve this goal.

Dynamic treatment regimes are routinely assigned to subjects participating in randomized medical trials when there is danger of serious side-effects necessitating cessation of treatment or when the optimal dosage at a given time should depend on the subjects clinical status or laboratory values. For example, in the active treatment arm of the Systolic Hypertension in the Elderly Program randomized trial (Cooperative Research Group, 1988; Borhani et al., 1991), subjects were assigned to a dynamic treatment regime that was intended to minimize the amount of medication required to maintain a subject’s systolic blood pressure (SBP) at or below a pre-determined goal. The treatment schedule was quite complex. Initially, all treated subjects were assigned the same dosage of anti-hypertensive medication; at eight weeks, subjects whose SBP exceeded the goal had their dosage doubled; at sixteen weeks, subjects whose SBP still exceeded the goal had a second anti-hypertensive added. At twenty-four weeks, a subject whose SBP continued to exceed the goal had their dose of the second anti-hypertensive doubled. The treatment schedule specified that subjects with evidence of severe clinical or laboratory toxicity would first have their dose of anti-hypertensive medications reduced; if the toxicity persisted, they would be switched to

an alternate medication. As in the Fast Track program, this dynamic treatment is used to tailor the dosage to the subject. Furthermore the use of rules for reducing dosage or switching medications due to side effects is a useful aid in reducing unexplained noncompliance. Indeed subjects whose medication is switched in accordance with the rules are compliers.

In this paper we consider methodology for estimation of the mean response to both nonrandom and random dynamic treatment regimes when the available data are observational. In a nonrandom dynamic treatment regime, the rules specifying the treatment level, as outputs of present need, are nonrandom. Both the Fast Track program and the Systolic Hypertension in the Elderly Program use nonrandom dynamic treatment regimes. In a random dynamic treatment regime, the time-specific treatment level is drawn from a conditional probability distribution depending only on measures of present needs. In Section 3 we provide quantitative definitions of these two types of dynamic treatment regimes. The modeling of responses to dynamic treatment regimes based on observational data has received very little statistical attention outside of a series of papers by Robins (1986, 1989, 1993, 1997). In these papers Robins considers the use of nested structural models to estimate a variety of conditional treatment effect parameters in the distribution of the potential response. Also Robins (1993) discusses estimating parameters in the response to a nonrandom dynamic treatment regime by censoring the subject at the first time the subject's treatment differs from the treatment as specified by the dynamic regime.

In Section 2 we review the underlying causal theory based on counterfactual or potential outcomes. Next in section 3 we discuss the assumption of sequential randomization along with the definitions of nonrandom and random dynamic treatment regimes. Section 4 uses the work in the previous sections to precisely specify the estimand in terms of the observational data distribution. We provide an estimating function in Section 5 and then in the last section we return to the Fast Track example.

2. POTENTIAL OUTCOMES

We use counterfactual or potential outcome models to quantify the desired treatment effect and to state assumptions. Neyman (1923) introduced counterfactual outcomes to analyze the causal effect of time-independent treatments in randomized studies. Rubin (1978) explicated Neyman’s ideas and extended Neyman’s work to the analysis of causal effects of time-independent treatments from observational data. Robins (1986, 1987) proposed a formal theory of causal inference that extended both Neyman’s and Rubin’s work to assess the direct and indirect effects of time varying treatments from experimental and observational longitudinal studies. We use these works to first specify our observations in a unified way regardless of the manner in which treatment is selected/assigned.

Suppose that the treatment lasts for K intervals; during this time period intermediate outcomes may be measured and at the end of the K intervals a response is measured. We denote the treatment regime/vector across the K intervals by $\bar{a}_K = (a_1, a_2, \dots, a_K)$; a_t is the level of treatment in the t th interval. In general we use a bar over a variable to denote that variable and all past values of the same variable, so $\bar{a}_t = (a_1, \dots, a_t)$. Let \mathcal{A}_K be the collection of all possible treatment vectors. Corresponding to each fixed value of the treatment vector, \bar{a}_K we conceptualize a potential (counterfactual) response denoted by $Y(\bar{a}_K)$ where $Y(\bar{a}_K)$ is the response at the end of the K th interval that a subject would have had if he/she followed the treatment regime, \bar{a}_K . The set of all potential responses is $\{Y(\bar{a}_K): \bar{a}_K \text{ varying in } \mathcal{A}_K\}$. This statement implicitly assumes that there is no need to index Y by others’ treatment or by the mechanism by which the treatment was selected; this is the Stable Unit Treatment Assumption (SUTVA, see Rubin, 1986). In the time varying treatment setting this is the consistency assumption of Robins (1997). In particular we assume that treatment of a subject does not influence any of the outcomes of any other subject (Cox, 1958); indeed we will make the stronger assumption that the observations on the sample of subjects are independent draws from one distribution. This assumption allows us to conceptualize a single potential outcome/response corresponding to each possible $\bar{a}_K \in \mathcal{A}_K$ and thus permit a well-defined, simple notation for an intermediate outcome and final response (see Robins, 1986 and for time-independent treatment, Rubin, 1986; Angrist, Imbens and Rubin, 1996).

SUTVA may well be violated in the Fast Track data. In addition to the home visiting, the Fast Track intervention included “friendship groups” in which small numbers of the intervention children were brought together in order to improve social skills. Thus in close friendships, the effect of home visiting on one child may affect the friend’s response. Alternately whether or not a dominant child is receiving home visits may alter the dynamics of the friendship groups and thus alter other children’s response to treatment. We do not deal with these complications here as an appropriate generalization of the proposed methodology would distract from the main points of this paper.

Capital letters are used to denote random variables. Treatment selection/assignment

is allowed to be stochastic and are denoted by the random vector, \bar{A}_K . So \bar{A}_K takes values in \mathcal{A}_K . Denote the variables available at the beginning of time interval t ($t > 0$) by $L_t = (S_t, V_t)$ where S_t is the severity vector and V_t is a vector of auxiliary information. In general both V_t and S_t may be summaries of information that are available prior to and at the beginning of time interval t . (S_t, V_t) is an intermediate outcome of (past) treatment. Denote the initial, time 0, variables by $L_0 = (Z, S_0, V_0)$ where Z consists of variables defining interesting subpopulations and S_0, V_0 have the same interpretation as above. Additionally in randomized studies Z may contain the indicator of the treatment regime to which a study subject was assigned. Corresponding to each treatment vector, \bar{a}_K , SUTVA implies that the intermediate outcome at time t can be written unambiguously as $(S_t(\bar{a}_t), V_t(\bar{a}_t))$ (see Robins,1997) and the potential response as $Y(\bar{a}_K)$.

A subject's potential outcomes, that is the "complete" severity and response data are $O_{sr} = \{Z, S_0, S_1(a_1), \dots, S_{K-1}(\bar{a}_{K-1}), Y(\bar{a}_K); \bar{a}_K \in \mathcal{A}_K\}$ and the complete auxiliary data are $O_{aux} = \{V_0, V_1(a_1), \dots, V_{K-1}(\bar{a}_{K-1}); \bar{a}_K \in \mathcal{A}_K\}$. Assuming SUTVA we may write the observed data in a unified way regardless of the manner in which treatment is selected/assigned. In both randomized and observational studies, the observed data are the pretreatment information plus the potential outcomes corresponding to the treatment pattern \bar{A}_K . That is, we observe $X = \{Z, S_0, V_0, A_1, S_1(A_1), V_1(A_1), A_2, \dots, S_{K-1}(\bar{A}_{K-1}), V_{K-1}(\bar{A}_{K-1}), A_K, Y(\bar{A}_K)\}$. In the following we use either Y or $Y(\bar{A}_K)$ to denote the observed response at the end of time interval K , either S_t or $S_t(\bar{A}_t)$ to denote the observed severity at the beginning of time interval t and either V_t or $V_t(\bar{A}_t)$ to denote the observed auxiliary variables available at the beginning of time interval t . Thus we may denote an individual's data in order of observation by

$$X = \{(Z, S_0, V_0), A_1, (S_1, V_1), A_2, \dots, (S_{K-1}, V_{K-1}), A_K, Y\}.$$

3. DYNAMIC REGIMES AND SEQUENTIAL RANDOMIZATION

In a dynamic treatment regime, the rule for treatment assignment in each of the K intervals may be random or nonrandom; in either case the rule depends only on prior severity. If the rules are nonrandom, denote the regime by the K vector, $\bar{d}_K = (d_1, \dots, d_K)$; at time t , treatment assignment is, $A_t = d_t(S_{t-1})$. (The Fast Track example uses nonrandom rules.) If the rules are random (stochastic), the treatment regime is characterized by a K vector of stochastic rules or equivalently treatment assignment probabilities, say $\bar{p}_K = (p_1, \dots, p_K)$; at time t , treatment assignment is drawn from the conditional distribution, $p_t(\cdot | S_{t-1})$. We shall denote by $P_{\bar{p}_K}$, the distribution of $(O_{sr}, O_{aux}, \bar{A}_K)$ had the entire study population been subjected to the dynamic regime \bar{p}_K . The associated expectation is denoted by $E_{\bar{p}_K}$. Note that a nonrandom dynamic treatment regime, \bar{d}_K , is just a special case of a random dynamic treatment regime with $p_t(a | S_{t-1})$ degenerate at the point, $a = d_t(S_{t-1})$. Henceforth, we do not, except where necessary, distinguish in our notation between random and nonrandom dynamic treatment regimes.

Dynamic treatment regimes should not be confused with nondynamic treatment regimes. In a nondynamic treatment regime, the variation in treatment level across time and subjects is specified prior to treatment onset. In other words, a nondynamic treatment regime is a special case of a dynamic treatment regime in which the treatment assignments do not vary by post-treatment observations (i.e. the rule, d_t is not a function of post-treatment information in S_{t-1} and is at most a function of pretreatment variables, L_0). Concrete examples of nondynamic treatment regimes are “assign 16 home visits in every semester following kindergarten,” or “assign 16 home visits in the first semester following kindergarten and 8 home visits in each semester thereafter.” Regression analyses that model the mean of $Y(\bar{a}_K)$ as \bar{a}_K varies are regression analyses comparing different nondynamic treatment regimes, each corresponding to a different treatment pattern, \bar{a}_K . For an example of this type of analysis see Robins et al. (1998) and Hernán et al. (1998).

Denote the distribution of $(O_{sr}, O_{aux}, \bar{A}_K)$ in our observational study by P_{obs} and the expectation with respect to this distribution by E_{obs} . Note that the marginal distribution of the complete data (O_{sr}, O_{aux}) is the same for both distributions, $P_{\bar{p}_K}$ and P_{obs} . Only the marginal/conditional distribution of the treatment (\bar{A}_K) varies between our observational distribution and the distribution $P_{\bar{p}_K}$. Thus we denote the marginal distribution of (O_{sr}, O_{aux}) by P without a subscript and expectation with respect to this marginal distribution by E without a subscript.

If the study population follows the dynamic treatment regime, \bar{p}_K , then A_t is conditionally independent of the potential outcomes, (O_{sr}, O_{aux}) , conditional on S_{t-1} . However in our observational study, A_t may not be conditionally independent of the potential outcomes, (O_{sr}, O_{aux}) , conditional on S_{t-1} , because there may be unmeasured “confounders” that determine treatment and are associated with the potential outcomes. In general, assumptions, in addition to SUTVA, about this distributional relationship (based on substantive knowl-

edge) must be used to identify causal effects and thus permit causal inference (for discussion, see section 11, of Robins, 1997). In this paper we use additional confounders in place of the unmeasured confounders and we assume:

Sequential Randomization: For each $t = 1, \dots, K$, A_t is independent of O_{sr} given $\{L_0, A_1, L_1, A_2, \dots, L_{t-1}\}$.

That is, the auxiliary data, the V 's, are sufficiently rich so the observational distribution satisfies the above sequential randomization. This means that within levels of $L_0, A_1, L_1, A_2, \dots, L_{t-1}$, we assume that the subjects with different levels of treatment do not vary systematically by the potential severity measures or responses. See Robins (1997) for further discussion of sequential randomization.

We have chosen to assume sequential randomization of the treatment level, \bar{A}_K with respect to O_{sr} only and not with respect to both (O_{sr}, O_{aux}) . There are two reasons for this. First there are practical, real-life examples in which sequential randomization of \bar{A}_K with respect to O_{sr} may be plausible but sequential randomization of \bar{A}_K with respect to O_{aux} is clearly untrue (see example 1 of Robins, 1987). The second reason is more philosophical. If a time varying correlate of both treatment level selection and response becomes available and the above sequential randomization assumption is more plausible when the V 's include this time varying covariate, then we would like to include this covariate in the V 's without having to make the additional assumption that the treatment levels are sequentially randomized with respect to this covariate as well.

4. IDENTIFICATION OF THE ESTIMAND IN TERMS OF THE OBSERVATIONAL DATA

Recall that the goal of this paper is to estimate, using observational data, the mean response that would have been observed had, contrary to fact, the entire study population followed a particular dynamic regime with rules \bar{p}_K . Thus we wish to estimate $E_{\bar{p}_K}[Y|Z]$ for Z denoting an interesting subpopulation. However we do not have data from a random sample of individuals following the dynamic regime, \bar{p}_K . Rather the available data are observational (from the P_{obs} distribution). The following lemma allows us to write $E_{\bar{p}_K}[Y|Z]$ in terms of the observational distribution, thus demonstrating that $E_{\bar{p}_K}[Y|Z]$ is identifiable from the available data. The lemma achieves this by providing a Radon-Nikodym derivative connecting the distribution $P_{\bar{p}_K}$ resulting from implementing the dynamic regime to the observational distribution P_{obs} . A secondary consequence of the lemma is to provide a way to construct estimating equations for $E_{\bar{p}_K}[Y|Z]$.

Let $\pi_t(\cdot|\bar{a}_{t-1}, \bar{l}_{t-1})$ represent the conditional probability mass function of A_t given $\bar{A}_{t-1} = \bar{a}_{t-1}, \bar{L}_{t-1} = \bar{l}_{t-1}$ in the observational distribution. Define

$$B(o_{sr}, \bar{a}_K) = P_{obs} \left[\prod_{t=1}^K \pi_t(a_t|\bar{a}_{t-1}, \bar{L}_{t-1}(\bar{a}_{t-1})) > 0 \mid O_{sr} = o_{sr} \right]$$

and

$$W_{\bar{p}_j}(\bar{a}_j, \bar{l}_{j-1}) = \prod_{t=1}^j \frac{p_t(a_t | s_{t-1})}{\pi_t(a_t | \bar{a}_{t-1}, \bar{l}_{t-1})}$$

for any $j \leq K$.

Lemma 4.1. Assume that the observational distribution, P_{obs} satisfies the sequential randomization assumption. If

$$P_{\bar{p}_K} [B(O_{sr}, \bar{A}_K) = 1] = 1 \quad (4.1)$$

then the distribution of $(Y, \bar{S}_{K-1}, \bar{A}_K, Z)$ under $P_{\bar{p}_K}$ is absolutely continuous with respect to the distribution of $(Y, \bar{S}_{K-1}, \bar{A}_K, Z)$ under P_{obs} and a version of the Radon-Nikodym derivative is

$$E_{obs} \left[W_{\bar{p}_K}(\bar{A}_K, \bar{L}_{K-1}) \middle| Y = y, \bar{S}_{K-1} = \bar{s}_{K-1}, \bar{A}_K = \bar{a}_K, Z = z \right]. \quad (4.2)$$

See the appendix for a proof of this lemma.

An intuitive interpretation of the lemma's assumption (4.1) is that, "Any treatment pattern that can result in the implementation of the dynamic treatment regime, \bar{p}_k , must also have a positive probability of occurring in the observational study." This assumption is eminently sensible since if a particular treatment pattern cannot occur in the observational study, then inference involving the response to this treatment pattern requires further knowledge/assumptions.

As stated above there are two important consequences of the fact that (4.2) is a Radon-Nikodym derivative. Using the lemma we have that the estimand can be written in terms of the observational distribution as

$$\begin{aligned} E_{\bar{p}_K} [Y | Z] &= E_{obs} \left[E_{obs} [W_{\bar{p}_K}(\bar{A}_K, \bar{L}_{K-1}) | Y, \bar{S}_{K-1}, \bar{A}_K, Z] Y \middle| Z \right] \\ &= E_{obs} \left[W_{\bar{p}_K}(\bar{A}_K, \bar{L}_{K-1}) Y \middle| Z \right]. \end{aligned} \quad (4.3)$$

More explicitly, the right hand side is, in terms of conditional densities, $E_{\bar{p}_K} [Y | Z = z] =$

$$\sum_{\bar{a}_K \in \mathcal{A}} \int_{\bar{l}_{K-1}} \int_y y f_Y(y | \bar{a}_K, \bar{v}_{K-1}, \bar{s}_{K-1}) \prod_{t=1}^K p_t(a_t | s_{t-1}) \prod_{t=1}^{K-1} f_t(l_t | \bar{a}_t, \bar{l}_{t-1}) f_0(l_0 | z) dy d\bar{l}_{K-1} \quad (4.4)$$

where f_Y, f_K, \dots, f_0 denote conditional densities for the observational distribution. The conditional density of Y given \bar{A}_K, \bar{L}_{K-1} is f_Y and the conditional density of L_t given $\bar{A}_{t-1}, \bar{L}_{t-1}$ is f_t . Thus the conditional mean is a function of the observational distribution via the conditional densities f_Y and f_{K-1}, \dots, f_0 . This is Robins (1986) G-computation formula for a dynamic regime.

The second consequence of the fact that (4.2) is a Radon-Nikodym derivative is that given any $P_{\bar{p}_K}$ -unbiased estimating equation for parameters in the distribution, $P_{\bar{p}_K}$ we can

weight this estimating equation by the Radon-Nikodym derivative and thereby produce an unbiased estimating equation distribution for the same parameters but now using data from the observational study, the P_{obs} distribution. Suppose for the moment that $\bar{\pi}_K$ is known. Let $\mu(\beta, Z)$ be a parameterization of $E_{\bar{p}_K}[Y|Z]$ where β is an unknown vector parameter and μ is a known function. Given data from $P_{\bar{p}_K}$, a simple estimating function for β , might be based on $\dot{\mu}(\beta, Z)(Y - \mu(\beta, Z))$ where $\dot{\mu}(\beta, Z)$ is the derivative of μ with respect to β . Given data from the P_{obs} distribution, we weight to produce the unbiased estimating function,

$$\mathbb{P}_n [W_{\bar{p}_K}(\bar{A}_K, \bar{L}_K)\dot{\mu}(\beta, Z)(Y - \mu(\beta, Z))] \quad (4.5)$$

where for a function g , $\mathbb{P}_n g(X)$ is defined as the average of g over the observational sample ($\mathbb{P}_n g(X) = 1/n \sum_{i=1}^n g(X_i)$).

Note that the denominator of $W_{\bar{p}_K}$ is informally “the probability of having the treatment pattern one did indeed have” in the observational study. Robins (1998) introduced the idea of weighting estimating functions by the inverse of this probability in order to estimate parameters of marginal structural models and direct effect structural nested models for time varying treatments. Robins refers to such estimators as inverse probability of treatment weighted estimators Robins (1998, 1999).

In the special case in which \bar{p}_K represents a nonrandom dynamic treatment regime, an alternate strategy is to use the inverse probability of censoring weighted estimators as considered by Robins and Rotnitzky (1992) and Rotnitzky and Robins (1995). Instead of modeling the treatment selection given past information, they model the adherence/non-adherence to the treatment regime given past information. Subjects are censored when their treatment first deviates from the dynamic treatment regime; these subjects receive a weight of zero. Subjects whose treatment patterns match the dynamic treatment regime receive a weight that is the product of the modelled adherence probabilities. Note that in this paper, for each t , we could separate the selection probability, $\pi_t(a_t|\bar{a}_{t-1}, \bar{l}_{t-1})$ into the product of the probability of adherence/non-adherence to the treatment regime given $\bar{A}_{t-1} = \bar{a}_{t-1}$, $\bar{L}_{t-1} = \bar{l}_{t-1}$ times the probability of $A_t = a_t$ given adherence or non-adherence to the treatment regime and $\bar{A}_{t-1} = \bar{a}_{t-1}$, $\bar{L}_{t-1} = \bar{l}_{t-1}$.

If the observational distribution satisfies sequential randomization then $\mu(\beta, Z) = E_{\bar{p}_K}[Y|Z]$ and β can be interpreted as a parameter from the $P_{\bar{p}_K}$ distribution. That is the sequential randomization assumption allows us to interpret $\mu(\beta, Z)$ as the mean response to the regime \bar{p}_K . Alternately we may view $\mu(\beta, Z)$ simply as a parameterization of (4.4); this viewpoint does not require sequential randomization and does not involve consideration of potential outcomes. We use this latter viewpoint in forming the estimating equation and in considering efficiency improvements.

5. AN ESTIMATING EQUATION

Recall that we wish to estimate $E_{\bar{p}_K}[Y|Z]$ for Z denoting an interesting subpopulation and the available data are observational (from the P_{obs} distribution). Assuming SUTVA, (4.1) and sequential randomization, $E_{\bar{p}_K}[Y|Z]$ is equal to (4.4) or equivalently (4.3). We separate this interpretation from the estimation. In this section we simply consider a parameterization of (4.4) denoted by $\mu(\beta, Z)$. In order to estimate β we no longer need SUTVA, sequential randomization or (4.1); indeed we no longer need to assume the existence of potential outcomes. However we assume regularity conditions for the asymptotic theory and that

$$E_{obs}W_{\bar{p}_K}(\bar{A}_K, \bar{L}_{K-1}) = 1. \quad (5.1)$$

This is easily shown to be a consequence of (4.1); however, (5.1) makes no reference to potential outcomes.

Note that $\mu(\beta, Z)$ is a function of the conditional densities f_Y and f_{K-1}, \dots, f_0 . On the other hand, the likelihood of the observational data can be written as a product of conditional densities:

$$\left(f_Y(y|\bar{a}_K, \bar{l}_{K-1}) \prod_{t=1}^{K-1} f_t(l_t|\bar{a}_t, \bar{l}_{t-1}) f_0(l_0|z) \right) \times \prod_{t=1}^K \pi_t(a_t|\bar{a}_{t-1}, \bar{l}_{t-1}), \quad (5.2)$$

where $\pi_t(\cdot|\bar{a}_{t-1}, \bar{l}_{t-1})$ represents the conditional probability mass function of A_t given $\bar{A}_{t-1} = \bar{a}_{t-1}, \bar{L}_{t-1} = \bar{l}_{t-1}$ in the observational distribution. Thus the estimand, (4.4), depends only on the first term in the likelihood. The second term in the likelihood concerns only the treatment selection probabilities ($\bar{\pi}_K$) which are nuisance parameters. An optimal estimating function (one resulting in an efficient estimator of β) will be orthogonal to the score functions for the nuisance parameters (Bickel et al., 1993). This means we have the potential to improve estimators based on (4.5) by replacing (4.5) by itself minus its projection on the score functions for the treatment selection probabilities (Robins, 1999). We do this in the following. Recall the observed data for one person is denoted by X . Set $U(X, \beta)$ equal to $W_{\bar{p}_K}(\bar{A}_K, \bar{L}_K)\dot{\mu}(\beta, Z)(Y - \mu(\beta, Z))$. The estimating function, (4.5) is just $\mathbb{P}_n U(X, \beta)$. The projection of $U(X, \beta)$ on the score functions of the treatment selection probabilities is given by $\sum_{t=1}^K E_{obs}[U(X, \beta)|\bar{L}_{t-1}, \bar{A}_t] - E_{obs}[U(X, \beta)|\bar{L}_{t-1}, \bar{A}_{t-1}]$ (Robins, 1999).

After algebraic simplification we have that

$$\begin{aligned} U(X, \beta) - \sum_{t=1}^K E_{obs}[U(X, \beta)|\bar{L}_{t-1}, \bar{A}_t] + E_{obs}[U(X, \beta)|\bar{L}_{t-1}, \bar{A}_{t-1}] = \\ W_{\bar{p}_K}(\bar{A}_K, \bar{L}_{K-1})\dot{\mu}(\beta, Z)(Y - \mu(\beta, Z)) - \sum_{t=1}^K W_{\bar{p}_t}(\bar{A}_t, \bar{L}_{t-1})\dot{\mu}(\beta, Z)(g_t(\bar{A}_t, \bar{L}_{t-1}) - \mu(\beta, Z)) \end{aligned}$$

$$+ \sum_{t=1}^K \sum_{a_t} \pi_t(a_t | \bar{A}_{t-1}, \bar{L}_{t-1}) W_{\bar{p}_t}(a_t, \bar{A}_{t-1}, \bar{L}_{t-1}) \dot{\mu}(\beta, Z)(g_t(a_t, \bar{A}_{t-1}, \bar{L}_{t-1}) - \mu(\beta, Z)) \quad (5.3)$$

where $W_{\bar{p}_t}(\bar{a}_t, \bar{l}_{t-1})$ represents $\prod_{u=1}^t \frac{p_u(a_u | s_{u-1})}{\pi_u(a_u | \bar{a}_{u-1}, \bar{l}_{u-1})}$ and

$$\begin{aligned} g_K(\bar{A}_K, \bar{L}_{K-1}) &= E_{obs}(Y | \bar{A}_K, \bar{L}_{K-1}) \\ g_t(\bar{A}_t, \bar{L}_{t-1}) &= E_{obs} \left(\sum_{a_{t+1}} p_{t+1}(a_{t+1} | S_t) g_{t+1}(a_{t+1}, \bar{A}_t, \bar{L}_t) \middle| \bar{A}_t, \bar{L}_{t-1} \right) \end{aligned} \quad (5.4)$$

for $t = 1, \dots, K-1$. If the parameterization of $\mu(\beta, Z)$ is saturated then the right hand side of (5.3) is the efficient influence function for β . See the appendix for proofs.

Note that whereas the treatment selection probabilities for the $P_{\bar{p}_K}$ are known – they are \bar{p}_K – in our observational study the treatment selection probabilities, $\bar{\pi}_K$, are unknown and thus must be estimated from the observational data. Additionally we will need to estimate $\{g_t(\bar{a}_t, \bar{l}_{t-1}), t = 1, \dots, K\}$. Given a model for the treatment selection probabilities, say $\pi_t(a_t | \bar{a}_{t-1}, \bar{l}_{t-1}; \alpha_w), t = 1, \dots, K$ with parameter, α_w , we can use maximum likelihood based on (5.2) to estimate α_w . We maximize,

$$\mathbb{P}_n \sum_{t=1}^K \log \pi_t(A_t | \bar{A}_{t-1}, \bar{L}_{t-1}; \alpha_w) \quad (5.5)$$

to get $\hat{\alpha}_w$ and then by substitution the vector, $\{\hat{\pi}_t, t = 1, \dots, K\}$. Next we parameterize $\{g_t(\bar{a}_t, \bar{l}_{t-1}), t = 1, \dots, K\}$ with the vector parameter, α_g and denote the parameterization by $\{g_t(\bar{a}_t, \bar{l}_{t-1}; \alpha_g), t = 1, \dots, K\}$. A variety of methods may be used to estimate α_g ; we use the following estimating equation, i.e., solve,

$$\begin{aligned} 0 &= \mathbb{P}_n (Y - g_K(\bar{A}_K, \bar{L}_{K-1}; \alpha_g)) \frac{\partial}{\partial \alpha_g} g_K(\bar{A}_K, \bar{L}_{K-1}; \alpha_g) + \\ &\sum_{t=1}^{K-1} \mathbb{P}_n \left(\sum_{a_{t+1}} p_{t+1}(a_{t+1} | S_t) g_{t+1}(a_{t+1}, \bar{A}_t, \bar{L}_t; \alpha_g) - g_t(\bar{A}_t, \bar{L}_{t-1}; \alpha_g) \right) \frac{\partial}{\partial \alpha_g} g_t(\bar{A}_t, \bar{L}_{t-1}; \alpha_g) \end{aligned} \quad (5.6)$$

to get $\hat{\alpha}_g$.

To use (5.3) for estimation of β , we substitute estimators of the parameters, α_w and α_g in the weights and in the g_t 's and set,

$$\begin{aligned} &\mathbb{P}_n W_{\bar{p}_K}(\bar{A}_K, \bar{L}_{K-1}; \hat{\alpha}_w) \dot{\mu}(\beta, Z)(Y - \mu(\beta, Z)) \\ &- \mathbb{P}_n \sum_{t=1}^K W_{\bar{p}_t}(\bar{A}_t, \bar{L}_{t-1}; \hat{\alpha}_w) \dot{\mu}(\beta, Z)(g_t(\bar{A}_t, \bar{L}_{t-1}; \hat{\alpha}_g) - \mu(\beta, Z)) \\ &+ \mathbb{P}_n \sum_{t=1}^K \sum_{a_t} \pi_t(a_t | \bar{A}_{t-1}, \bar{L}_{t-1}; \hat{\alpha}_w) W_{\bar{p}_t}(a_t, \bar{A}_{t-1}, \bar{L}_{t-1}; \hat{\alpha}_w) \dot{\mu}(\beta, Z) \end{aligned}$$

$$(g_t(a_t, \bar{A}_{t-1}, \bar{L}_{t-1}; \hat{\alpha}_g) - \mu(\beta, Z)), \quad (5.7)$$

to 0 and solve for β .

Under regularity conditions, the following two properties hold.

(1) First assuming the model for μ is correct, and if the model for $\bar{\pi}_K$ OR $\{g_t(\bar{a}_t, \bar{l}_{t-1}), t = 1, \dots, K\}$ is correct, the resulting estimator of β is consistent and asymptotically normal.

(2) The asymptotic variance of $\hat{\beta}$ can be estimated by a sandwich estimator. Define $\hat{\Sigma} =$

$$\mathbb{P}_n \left(S_\beta(X; \hat{\alpha}_w, \hat{\alpha}_g, \hat{\beta}) + \hat{S}_{\beta,w} \hat{I}_w^{-1} S_w(X; \hat{\alpha}_w) + \hat{S}_{\beta,g} \hat{I}_g^{-1} S_g(X; \hat{\alpha}_g) \right)^{\otimes 2}$$

where for vector V , $V^{\otimes 2} = VV^T$, $S_\beta(X; \hat{\alpha}_w, \hat{\alpha}_g, \beta)$ is the estimating function for β (formula (5.7) is $\mathbb{P}_n S_\beta$), $S_w(X; \alpha_w)$ is the score function for α_w (the derivative of (5.5) with respect to α_w is $\mathbb{P}_n S_w(X; \alpha_w)$), \hat{I}_w is the observed information matrix for α_w (minus the Hessian of (5.5)), $S_g(X; \alpha_g)$ is the estimating function for α_g (formula (5.6) is $\mathbb{P}_n S_g$), \hat{I}_g is minus the derivative of (5.6) with respect to α_g , $\hat{S}_{\beta,w} = \mathbb{P}_n \frac{\partial}{\partial \alpha_w} S_\beta(X; \hat{\alpha}_w, \hat{\alpha}_g, \hat{\beta})$ and $\hat{S}_{\beta,g} = \mathbb{P}_n \frac{\partial}{\partial \alpha_g} S_\beta(X; \hat{\alpha}_w, \hat{\alpha}_g, \hat{\beta})$. Also define $\hat{B} = \mathbb{P}_n \dot{\mu}(\beta, Z) \dot{\mu}(\beta, Z)^T$. Then a consistent estimator of the asymptotic variance of $\hat{\beta}$ is given by $\hat{B}^{-1} \hat{\Sigma} \hat{B}^{-1}$.

See the appendix for proofs of these results.

Consistent models for g_1, \dots, g_K : The relationships among the g_1, \dots, g_K are highly constrained. Thus it is difficult to formulate models for g_1, \dots, g_K that are consistent with one another. For example, models for g_K and g_{K-1} are constrained by the following relationships,

$$g_{K-1}(\bar{a}_{K-1}, \bar{l}_{K-2}) = \int_{l_{K-1}} \sum_{a_K} p_K(a_K | s_{K-1}) g_K(\bar{a}_K, \bar{l}_{K-1}) f_{K-1}(l_{K-1} | \bar{a}_{K-1}, \bar{l}_{K-2}) dl_{K-1}$$

where p_K is specified by the treatment regime and f_{K-1} is the unspecified conditional density of L_{K-1} given $(\bar{A}_{K-1}, \bar{L}_{K-2})$. Or equivalently, given models for g_K and g_{K-1} there must be exist a conditional density f_{K-1} for which the above display is true. One way to formulate g_1, \dots, g_K which satisfy the above interrelationships is to parameterize each g_t as linear in \bar{l}_{t-1} and independent of \bar{a}_t . This particular parameterization (combine above display and (5.8) below) is consistent with no treatment effect. If there is a treatment effect, modeling the g_t 's in this way is used to improve efficiency. We use this parameterization in our example. See Robins and Wasserman (1997) and Scharfstein et al. (1999) for discussions about consistent parametrization.

Double Robustness: Instead of using (5.7) to estimate β , we could use (4.5) with estimated weights. Consistency of the resulting $\hat{\beta}$ would require that $\bar{\pi}_K$ is correctly parameterized. Alternately we could base estimation of β on the formula in (4.4). Note that (4.4) can be written in terms of the g_t 's as

$$\mu(\beta, Z) = \sum_{a_1} E[g_1(a_1, L_0) p_1(a_1 | S_0) | Z]. \quad (5.8)$$

As noted previously, this is the G-computation formula of Robins (1986). Thus to directly estimate β using (4.4) we may estimate the g_t 's in order to arrive at an estimate of g_1 and then solve

$$0 = \mathbb{P}_n \dot{\mu}(\beta, Z) \left(\sum_{a_1} g_1(a_1, L_0; \hat{\alpha}_g) p_1(a_1 | S_0) - \mu(\beta, Z) \right). \quad (5.9)$$

We can only expect the resulting $\hat{\beta}$ to be a consistent estimator of β if we have correctly parameterized the g_t 's.

An advantage of the use of (4.5) over (5.9) is that it is easier to parameterize the $\bar{\pi}_K$ than it is to parameterize the \bar{g}_K due to the constrained relationships among the \bar{g}_K . An advantage of (5.7) over (5.9) and over the use of (4.5) is that the use of (5.7) to estimate μ leads to a consistent estimator if either the g_t 's or π_t 's are parameterized correctly (see (1) above). This is the “double robustness” property; see Scharfstein et al. (1999) and Robins (2000) for further discussion. Furthermore note that the first term in (5.7) is (4.5) with estimated weights and the $t = 1$ summand in the last term is (5.9). Thus (5.7) combines both the direct estimation method and the weighted estimation method (4.5) so as to achieve consistency even when only the g_t 's or only the π_t 's are parameterized correctly.

6. EXAMPLE

The Fast Track trial is a longitudinal, multi-site, multi-cohort trial of a preventive intervention versus a control. For the purposes of this illustration we use a subset of the children for whom data was available at the time of the analysis; this is 579 high risk children of which 202 are in the intervention group. This is not a representative subset; furthermore the variables used in the analysis were selected for illustrative purposes only. A more complete analysis is forthcoming. We consider two endpoints, specifically chosen to illustrate the diversity of results that can follow from the proposed analyses. Both endpoints are from end of grade 3 teacher evaluations. The first endpoint is a teacher rating of school behavior problems. The second endpoint is a teacher rating of behavior change over the course of grade 3. This second rating assesses improvement across the year in social and emotional adjustment.

Beginning in the fall of grade 2, a nonrandom dynamic treatment regime for home visiting was planned as part of the intervention. The home visits were designed to improve parental functioning. The time intervals are fall and spring of grade 2 and 3 ($K = 4$). The severity for semester t is S_t , the total score on the home visiting process measure; S_t is a measure, taking values in the interval $[1, 24]$, evaluating the quality of family functioning. High values indicate better functioning. The treatment rule is: if assigned to control, there is no assignment of home visiting by staff; if assigned to treatment the staff should follow the treatment rule,

$$d_t(S_{t-1}) = 16I\{S_{t-1} \leq 8\} + 8I\{9 \leq S_{t-1} \leq 16\} + 4I\{17 \leq S_{t-1}\}, t = 1, 2, 3, 4$$

in assigning home visiting level. Thus, if a child is in the intervention group and the total on the home visiting process measure from the end of grade 1, S_0 , is less than 8, staff were to assign 16 home visits for the fall of grade 2 and so on. This means that $A_t \in \{4, 8, 16\}$.

Our primary goal is an evaluation of the average effect of the nonrandom dynamic treatment regime on the two endpoints. This regime corresponds to degenerate treatment selection probabilities, i.e.,

$$p_t(a|s_{t-1}) = 1 \text{ if } a = d_t(s_{t-1}) \text{ and } 0, \text{ otherwise}$$

for $t = 1, 2, 3, 4$.

Staff were told that in exceptional cases they might need to deviate from the rule for recommending home visiting level; in practice, staff recommendations deviated from the rule on 22% of the 202 * 4 occasions. The staff recommended more home visits than the rule on 15% of the occasions and on 7% of the occasions the staff recommended fewer home visits. In all, 104 intervention children (52%) have treatment recommendation patterns that coincide with the rule for all 4 time periods. The staff did not/could not deviate from the rule for the control children as no home visiting assignments were made for the control children. Because the staff deviated from the rule(for the intervention children), we treat

the group of intervention children as an observational sample in which the home visiting recommendations may have been influenced by severity (S_{t-1}) in ways varying from the rule and indeed, the recommendations may have been influenced by measures other than severity. The Fast Track research team collected a vast amount of information relating to why staff recommended intervention levels. For the purposes of this example, we will use past recommended intervention level, the home visiting process measure and site and cohort indicators to model the distribution of home visiting recommendations (these will be included in the L 's). Also included in the L 's are time-varying predictors of the two endpoints (Recall that these variables are chosen for illustrative purposes; they do not encompass the full set of variables that might be used in a substantive treatment). See Table 1 for a description of L .

We assume that sequential randomization holds; that is we assume that within any cross-stratification of $\{L_0, A_1, L_1, A_2, \dots, L_{t-1}\}$ the staff treatment recommendation, A_t , is not predictive of the array of potential outcomes, O_{sr} . For example, this means that (within a cross-stratification) children recommended 16 home visits do not have a higher chance of high values of $\{Y(a_1, a_2, a_3, a_4), a_t = 4, 8, 16, t = 1, 2, 3, 4\}$ than children recommended 4 home visits. It also means that (within a cross-stratification) children recommended 16 home visits do not have higher chance of high values of $Y(16, 16, 16, 16) - Y(4, 4, 4, 4)$ than children recommended 4 home visits.

To implement (5.7) we first estimate the weights, $W_{\bar{p}_4}$. Since the staff were not to and did not assign any intervention to control children, the weights for the control children are equal to one. Furthermore, we do not need to estimate the treatment selection probabilities, $\bar{\pi}_K$, for the control children as we know that each probability has mass one at 0 home visiting. Consider the intervention group children. The numerator of $W_{\bar{p}_4}$ is one for an intervention group child whose treatment recommendations coincide with the treatment rule and zero otherwise. This means that of the 202 intervention children, 98 (202-104) will have $W_{\bar{p}_4} = 0$ and thus the first term in (5.7) will be zero for these children. This makes sense because without further (smoothness) assumptions the responses from the 98 children whose treatment recommendations are inconsistent with the regime are not informative of responses to the treatment regime. This issue is discussed in further detail below. To estimate the denominator of the weight, we fit a proportional odds model to the staff home visiting recommendations for the intervention group children. For each time t , we fit, via maximum likelihood,

$$\begin{aligned} \text{logit}(\pi_t(16|\bar{L}_{t-1}, \bar{A}_{t-1}; \alpha_w)) &= \alpha_{w,t01} + \alpha'_{w,1} ZZ(\bar{L}_{t-1}, \bar{A}_{t-1}) \\ \text{logit}(1 - \pi_t(4|\bar{L}_{t-1}, \bar{A}_{t-1}; \alpha_w)) &= \alpha_{w,t02} + \alpha'_{w,1} ZZ(\bar{L}_{t-1}, \bar{A}_{t-1}) \end{aligned}$$

where $\alpha_w = \{\alpha_{w,t01}, \alpha_{w,t02}, \alpha_{w,1}, t = 1, 2, 3, 4\}$ and $ZZ(\bar{L}_{t-1}, \bar{A}_{t-1})$ is a vector summary of past information. See Table 2 for the estimates of α_w . Note that in all four time periods, a low home visiting process measure score is predictive of higher staff home visiting

recommendation; also in the last three time periods, a past high staff home visiting recommendation is predictive of a higher staff home visiting recommendations. Additionally there are some minor differences in home visiting recommendations by cohort and site. Since there were only 4 intervals and the sample size was not low we choose to fit separate models for the home visiting recommendations in each interval; alternately we could have fit one model with possibly time varying intercepts and regression coefficients to all 4 intervals simultaneously (see Hernán et al. for an example, 1998a).

To implement (5.7) we must also estimate α_g . We estimate α_g separately for control and intervention groups. However since the last two terms in (5.7) cancel for control group children (because the weights are equal to one and the π_t have mass one for 0 home visits), we do not use the estimator of α_g for the control group. Thus we use only need to estimate α_g for the intervention group children. We model g_1 through g_4 by linear models:

$$g_t(\bar{A}_t, \bar{L}_{t-1}; \alpha_g) = \alpha_{g0} + \alpha'_{g1} XX(\bar{L}_{t-1})$$

where $XX(\bar{L}_{t-1})$ is a summary vector of past information and estimate $\alpha_g = \{\alpha_{g0}, \alpha_{g1}\}$ by solving (5.6). Note that the g_t 's are assumed to be functionally independent of both the present time t home visiting level and the past assigned home visiting levels (\bar{A}_t). It is easy to check that this assumption plus the linearity of g_t imply that the parameterizations of g_1, \dots, g_K will be consistent with one another. However as mentioned earlier this parameterization can only be correct under the null hypothesis of no treatment effect. Thus the use of this model for \bar{g}_K implies that we are assuming our model for the $\bar{\pi}_K$ is correct. The estimates of α_g are given in Table 3.

For each response, we model the mean response to the treatment regime, \bar{p}_4 by

$$\mu(\beta, Z, treatment) = \beta_0 + \beta_1 SBP1 + \beta_2 treatment$$

where *treatment* is an indicator with the value one for the intervention children and when grade3 teacher rating of school behavior problems is the response, grade 1 teacher rating of school behavior problems *SBP1* is included in the regression. When the teacher rating of behavior change over the course of grade 3 is the response, *SBP1* is not included. We estimate β by setting (5.7) to zero and solving for $\hat{\beta}$. See Table 4, $h = 1$, for estimates of β_2 . As a comparison we include the results of an “intent to treat” analysis (ITT), this is a simple linear regression of the response on treatment (and possibly pretreatment covariates). The ITT analysis does not adjust for the level of staff deviance from the rules of the treatment regime. In contrast the analysis of the dynamic treatment regime yields an estimate of the mean treatment effect in the setting in which staff follow the treatment regime rules in assigning home visiting. In Table 4a, we see that for the response, school behavior problems, both the ITT treatment effect and the treatment effect for the treatment regime ($h = 1$) are highly significant. But the estimated treatment effect in the scenario in which the staff follow the rules is nearly twice as large as the ITT treatment effect. Recall that (see

Table 2) in addition to the home visiting process measure, past treatment recommendation, site and cohort were predictive of the treatment recommendation probabilities ($\bar{\pi}_4$) in the observational data. The dynamic regime \bar{d}_4 forces uniformity across past treatment recommendation, site and cohort in making present treatment recommendations. It is unclear whether the increased treatment effect is due to this forced uniformity and/or due to using the home visiting measure only as prescribed by the regime rule. This is discussed further below. In contrast, in Table 4b, we see that while there is a significant ITT treatment effect on behavior change, there is no treatment effect for the treatment regime ($h = 1$). In this case it appears that staff judgment in deviating from the rules has enhanced the treatment effect. Similarly to before it is unclear whether the decreased treatment effect is due to forced uniformity across site and cohort in assigning treatment and/or due to ignoring past level of assigned treatment in making future recommendations and/or due to using the home visiting measure only as prescribed by the regime rule.

Recall that 98 of the 202 intervention children have $W_{\bar{p}_4} = 0$. These children's responses enter (5.7) only through the estimated g_t 's. In order to better utilize these responses and also obtain a better understanding of the reasons for the difference in treatment effects between the regime and the ITT analyses, we examine a small variety of random dynamic treatment regimes, denoted by $\bar{p}_{4,h}$, $h \in \{0, .2, .4, \dots, 1\}$. $p_{t,h}(a|s_{t-1})$ will be the probability of recommending home visiting level a at time t when the home visiting measure is equal to s_{t-1} . For each $t \in \{1, 2, 3, 4\}$, set $p_{t,h}$ equal to a mixture of the indicator, $I\{a = d_t(s_{t-1})\}$ and $q_t(a|s_{t-1})$;

$$p_{t,h}(a|s_{t-1}; \gamma) = h I\{a = d_t(s_{t-1})\} + (1 - h) q_t(a|s_{t-1}; \gamma)$$

where q_t is a conditional probability mass function for treatment selection given severity, s_{t-1} . We choose q_t to approximate the observational distribution of treatment recommendation at time t given only the home visiting process measure. All of these random dynamic regimes do not allow site, cohort and past home visiting recommendations to influence present home visiting recommendation. Note that $h = 1$ corresponds to originally planned treatment regime. The random dynamic treatment regime, $\{p_{1,h}, p_{2,h}, p_{3,h}, p_{4,h}\}$ corresponds to drawing a Bernoulli random variable with success probability equal to h at each time t ; if the Bernoulli outcome is one then set A_t equal to $d_t(S_{t-1})$, otherwise draw A_t from the probability mass function, $q_t(a|S_{t-1}; \gamma)$.

Of course, the observational distribution of treatment recommendations given only the home visiting process measure is unknown and must be approximated. We estimate this approximation. To construct q_t , $t = 1, 2, 3, 4$, at each time we fit, via maximum likelihood, the following proportional odds model to the data:

$$\begin{aligned} \text{logit}(q_t(16|S_{t-1}; \gamma)) &= \gamma_{t01} + \gamma_{t1} I\{d(S_{t-1}) = 16\} + \gamma_{t2} I\{d(S_{t-1}) = 4\} \\ \text{logit}(1 - q_t(4|S_{t-1}; \gamma)) &= \gamma_{t02} + \gamma_{t1} I\{d(S_{t-1}) = 16\} + \gamma_{t2} I\{d(S_{t-1}) = 4\} \end{aligned}$$

where $\gamma = \{\gamma_{t01}, \gamma_{t02}, \gamma_{t1}, \gamma_{t2}, t = 1, 2, 3, 4\}$. The estimated values of γ are given in Table 5. We use $q_t(\cdot|S_{t-1}; \hat{\gamma})$ to form the $p_{t,h}(\cdot|S_{t-1}; \hat{\gamma})$ when $h < 1$.

Note that because the proportional odds model, q_t is not a saturated model, the models for the q_t and π_t will not be consistent with one another. However we only view q_t as an approximation to the observational distribution of treatment recommendation given severity. Also note that the $p_{t,h}(\cdot|S_{t-1}; \hat{\gamma})$ are used to form the numerator of the weights and as before the $\pi_t(\cdot|\bar{L}_{t-1}, \bar{A}_{t-1}; \hat{\alpha}_w)$ are used to form the denominator of the weights. For each value of h and each of the two responses, we use (5.6) to construct estimators of α_g , but for $h < 1$, p_t is replaced by $p_{t,h}(\cdot|S_{t-1}; \hat{\gamma})$. The estimates of γ are omitted. The model for the mean treatment response is as before; Table 4 contains the results. The formula for the estimated standard errors changes since the numerator of the weights is now estimated. See the appendix for the appropriate formula. Consider the results for $h = 0$ in Table 4. These results are essentially equivalent to the ITT results. Recall that for $h = 0$, the dynamic treatment regime corresponds to drawing A_t from the probability mass function, $q_t(a|S_{t-1}; \gamma)$. Thus even though there was systematic variation from the treatment assignment rule, systematic variation that can be accounted for by past home visiting recommendation and site and cohort differences, this systematic variation does not appear to alter the mean response. Rather we see that as the random dynamic treatment regime approaches (h approaches 1) the original dynamic treatment regime, the mean effect changes. This indicates that the change in mean effect from the ITT analysis to the original dynamic treatment regime analysis is due to only using the home visiting measure as prescribed by the regime rule.

Clearly a more complete analysis of the Fast Track data is needed to investigate these substantive issues. As stated previously, the two responses were chosen specifically to illustrate the diversity of thought-provoking results that can be obtained by these types of analyses. Certainly one should give careful thought to which responses are of primary interest; variations in the dynamic treatment regime may differentially impact different responses.

7. DISCUSSION

This methodology provides a way to evaluate the effects of dynamic treatment regimes using observational data. It should be extended in a variety of ways. First the interpretation of the results depends heavily on the assumption of sequential randomization. It is difficult to believe that this assumption really holds in the observational setting. Thus an extension of the sensitivity analyses of Robins et al. (1999a) and Scharfstein et al. (1999) to this setting is needed. A second needed extension is to allow for missing severity measures. There are two ways to allow for missing severity measures. First, if one can expect severity measures to be occasionally missing in practice, then it makes sense to extend the dynamic regime rules to include rules for assignment when severity is missing. On the other hand if one wishes to make inference for a rule that does not allow for missing severity measures, yet one's observational data include individuals with missing severity measures then the methodology provided here must be adapted. One possibility is a modification of the weighting method as described in Hernán et al. (1998). A third extension would be to grouped data. Many intervention/prevention studies take place in a school based setting and thus grouped data dominates. Lastly more systematic work is needed on how to formulate/evaluate dynamic treatment rules, and in particular the evaluation and detection of which posttreatment conditions (such as the development of side effects or the occurrence of family crises) may negate or enhance the effect of treatment.

8. APPENDIX

Lemma 4.1

Proof.

Below we repeatedly use the notation, $Y == Y(\bar{A}_K)$, $L_t == (S_t, V_t)$ and $L_t == L_t(\bar{A}_t)$. Let U be a Borel set. We have that

$$\begin{aligned}
 & E_{obs} [I\{(Y, \bar{S}_{K-1}, \bar{A}_K, Z) \in U\} E_{obs} [W_{\bar{p}_K}(\bar{A}_K, \bar{L}_{K-1}) | Y, \bar{S}_{K-1}, \bar{A}_K, Z]] \\
 &= E_{obs} [I\{(Y, \bar{S}_{K-1}, \bar{A}_K, Z) \in U\} W_{\bar{p}_K}(\bar{A}_K, \bar{L}_{K-1})] \\
 &= \sum_{\bar{a}_K} E_{obs} [I\{(Y, \bar{S}_{K-1}, \bar{a}_K, Z) \in U\} W_{\bar{p}_K}(\bar{a}_K, \bar{L}_{K-1}) I\{\bar{A}_K = \bar{a}_K\}] \\
 &= \sum_{\bar{a}_K} E_{obs} \left[I\{(Y(\bar{a}_K), \bar{S}_{K-1}, \bar{a}_K, Z) \in U\} W_{\bar{p}_{K-1}}(\bar{a}_K, \bar{L}_{K-2}) p_K(a_K | S_{K-1}) \right. \\
 & \quad \left. P_{obs} [\pi_K(a_K | \bar{a}_{K-1}, \bar{L}_{K-1}) > 0 | O_{sr}, \bar{L}_{K-2}, \bar{A}_{K-1} = \bar{a}_{K-1}] I\{\bar{A}_{K-1} = \bar{a}_{K-1}\} \right].
 \end{aligned}$$

Continuing in this fashion, eliminating the indicator variables concerning A_j and accumulating the conditional expectations concerning the $\pi_j(a_j|\bar{L}_{j-1}(\bar{a}_{j-1}), \bar{a}_{j-1})$ we arrive at,

$$\sum_{\bar{a}_K} E_{obs} \left[I\{(Y(\bar{a}_K), \bar{S}_{K-1}(\bar{a}_{K-1}), \bar{a}_K, Z) \in U\} B_1(O_{sr}, \bar{a}_K) \prod_{t=1}^K p_t(a_t|S_{t-1}(\bar{a}_{t-1})) \right] \quad (8.1)$$

where B_1 is formed from repeated conditional expectations and can be defined iteratively beginning with,

$$B_K(O_{sr}, \bar{L}_{K-2}, \bar{a}_K) = P_{obs} \left[\prod_{t=1}^K \pi_t(a_t|\bar{a}_{t-1}, \bar{L}_{t-1}(\bar{a}_{t-1})) > 0 \middle| O_{sr}, \bar{L}_{K-2}, \bar{A}_{K-1} = \bar{a}_{K-1} \right]$$

and then for each $j \leq K-1$

$$B_j(O_{sr}, \bar{L}_{j-2}, \bar{a}_K) = E_{obs}[B_{j+1}(O_{sr}, \bar{L}_{j-1}, \bar{a}_K) | O_{sr}, \bar{L}_{j-2}, \bar{A}_{j-1} = \bar{a}_{j-1}]$$

and for $j=1$ as,

$$B_1(O_{sr}, \bar{a}_K) = E_{obs}[B_2(O_{sr}, L_0, \bar{a}_K) | O_{sr}].$$

Since the distribution of O_{sr} is the same under both $P_{\bar{p}_K}$ and P_{obs} distributions, (8.1) can also be written with $E_{\bar{p}_K}$ in place of E_{obs} . But then the sum over \bar{a}_K can be taken inside the expectation to yield

$$\begin{aligned} E_{obs} [I\{(Y, \bar{S}_{K-1}, \bar{a}_K, Z) \in U\} E_{obs} [W_{\bar{p}_K}(\bar{A}_K, \bar{L}_{K-1}) | Y, \bar{S}_{K-1}, \bar{A}_K]] = \\ E_{\bar{p}_K} [B_1(O_{sr}, \bar{A}_K) I\{(Y, \bar{S}_{K-1}, \bar{A}_K, Z) \in U\}]. \end{aligned}$$

Since B_1 can be expressed as repeated expectations of a probability (B_K) that is necessarily bounded above by one, (4.1) implies that $B_1(O_{sr}, \bar{A}_K) = 1$ *a.e.* Thus we have absolute continuity and (4.2) is the Radon-Nikodym derivative of $P_{\bar{p}_K}$ with respect to P_{obs} . ■

In the following we assume (5.1). Denote the true value of β by β_0 . In the following it is useful to note two alternate ways to rewrite (5.3). The first emphasizes the correct specification of $\bar{\pi}_K$:

$$\begin{aligned} S_\beta(X; \alpha_w, \alpha_g, \beta) &= W_{\bar{p}_K}(\bar{A}_K, \bar{L}_{K-1}) \dot{\mu}(\beta, Z) (Y - \mu(\beta, Z)) \\ &- \sum_{t=1}^K \sum_{a_t} (I\{A_t = a_t\} - \pi_t(a_t|\bar{A}_{t-1}, \bar{L}_{t-1})) W_{\bar{p}_t}(a_t, \bar{A}_{t-1}, \bar{L}_{t-1}) \dot{\mu}(\beta, Z) \\ &\quad (g_t(a_t, \bar{A}_{t-1}, \bar{L}_{t-1}) - \mu(\beta, Z)). \end{aligned} \quad (8.2)$$

Note that if $\bar{\pi}_K$ is correctly specified then the second term above has mean zero even if the g_t 's are incorrectly specified and/or $\beta \neq \beta_0$. The second way to rewrite (5.3) emphasizes the correct estimation of the g_t 's:

$$S_\beta(X; \alpha_w, \alpha_g, \beta) = W_{\bar{p}_K}(\bar{A}_K, \bar{L}_{K-1}) \dot{\mu}(\beta, Z) (Y - g_K(\bar{A}_K, \bar{L}_{K-1}))$$

$$\begin{aligned}
& - \sum_{t=1}^{K-1} W_{\bar{p}_t}(\bar{A}_t, \bar{L}_{t-1}) \dot{\mu}(\beta, Z) \left(g_t(\bar{A}_t, \bar{L}_{t-1}) - \sum_{a_{t+1}} p_{t+1}(a_{t+1}|S_t) g_{t+1}(a_t, \bar{A}_t, \bar{L}_t) \right) \\
& \quad + \dot{\mu}(\beta, Z) \left(\sum_{a_1} g_1(a_1, L_0) p_1(a_1|S_0) - \mu(\beta, Z) \right). \tag{8.3}
\end{aligned}$$

Note that if the g_t 's are correctly specified then the first two terms have expectation zero even if $\bar{\pi}_K$ is incorrectly specified and/or $\beta \neq \beta_0$.

Lemma 8.1. Formula (5.3) is orthogonal to, or equivalently, uncorrelated with the score functions for the treatment assignment probabilities, $\bar{\pi}_K$, thus (5.3) is $W_{\bar{p}_K}(\bar{A}_K, \bar{L}_K) \dot{\mu}(\beta, Z) (Y - \mu(\beta, Z))$ minus its projection on the score functions for the treatment assignment probabilities.

Proof. First note that all score functions for the $\bar{\pi}_K$ belong to the linear span of

$$B_t(a, \bar{A}_t, \bar{L}_{t-1}) = U_t(\bar{A}_{t-1}, \bar{L}_{t-1}) (I\{A_t = a\} - \pi_t(a|\bar{A}_{t-1}, \bar{L}_{t-1}))$$

for each possible treatment level, a and $t \in \{1, \dots, K\}$ and for $U_t(\bar{A}_{t-1}, \bar{L}_{t-1})$ for $t \in \{1, \dots, K\}$ arbitrary bounded functions. Thus we need only show that (5.3) or equivalently (8.2) is orthogonal to each of these functions. Note that $B_j(a, \bar{A}_j, \bar{L}_{j-1})$ is orthogonal to the terms with $t \neq j$ in the sum in (8.2). The covariance between $B_j(a, \bar{A}_j, \bar{L}_{j-1})$ and (8.2) is thus

$$E_{obs} [W_{\bar{p}_K}(\bar{A}_K, \bar{L}_{K-1}) \dot{\mu}(\beta, Z) (Y - \mu(\beta, Z)) B_j(a, \bar{A}_j, \bar{L}_{j-1})] \tag{8.4}$$

$$- E_{obs} \left(\sum_{a_j} (I\{A_j = a_j\} - \pi_j(a_j|\bar{A}_{j-1}, \bar{L}_{j-1})) \right)$$

$$W_{\bar{p}_j}(a_j, \bar{A}_{j-1}, \bar{L}_{j-1}) \dot{\mu}(\beta, Z) (g_j(a_j, \bar{A}_{j-1}, \bar{L}_{j-1}) - \mu(\beta, Z)) B_j(a, \bar{A}_j, \bar{L}_{j-1}). \tag{8.5}$$

Since given $\bar{A}_{j-1}, \bar{L}_{j-1}$, $B_j(a, \bar{A}_j, \bar{L}_{j-1})$ has mean zero, the second term (8.5) reduces to

$$- E_{obs} \sum_{a_j} I\{A_j = a_j\} W_{\bar{p}_j}(a_j, \bar{A}_{j-1}, \bar{L}_{j-1}) \dot{\mu}(\beta, Z) (g_j(a_j, \bar{A}_{j-1}, \bar{L}_{j-1}) - \mu(\beta, Z))$$

$$B_j(a, \bar{A}_j, \bar{L}_{j-1}).$$

But this is the negative of the first term (8.4) (repeatedly take conditional expectations and use (5.1)). ■

Lemma 8.2. If the parameterization of $\mu(\beta, Z)$ is saturated and g_1, \dots, g_K are nonparametric then (5.3) is the efficient influence function for β .

Proof. Because of the factorization, (5.2) of the likelihood and the fact that the functional $\mu(\beta, Z)$ does not depend on the factor $\prod_{t=1}^K \pi_t(a_t|\bar{a}_{t-1}, \bar{l}_{t-1})$, the efficient influence function

is the same in the model for which $\bar{\pi}_K$ is completely unknown as in the model where $\bar{\pi}_K$ is a known function up to a parameter, α_w (Robins et al., 1994; Robins and Ritov, 1997). However when $\mu(\beta, Z)$ is saturated, the model with $\bar{\pi}_K$ completely unknown is a nonparametric model for the observed data. Bickel et al. (1993) show that there is only one influence function for any nonparametric model, which therefore must necessarily be the efficient influence function. Thus if we find an influence function of the model in which $\bar{\pi}_K$ completely unknown, this will be the efficient influence function for this nonparametric model and also be the efficient influence function for the model in which $\bar{\pi}_K$ is parameterized. Now when $\bar{\pi}_K$ is completely known, $U(X, \beta, \bar{p}_K) = W_{\bar{p}_K}(\bar{A}_K, \bar{L}_K)\dot{\mu}(\beta, Z)(Y - \mu(\beta, Z))$ is an unbiased estimating function for β regardless of the distribution of $Y - \mu$. It follows that $U(X, \beta)$ minus its projection on all scores for $\bar{\pi}_K$ (as given in (5.3)) is orthogonal to the nuisance tangent space for the nonparametric models. Furthermore (5.3) is an influence function (the expectation of its derivative with respect to β is -1). Thus (5.3) is the efficient influence function when $\bar{\pi}_K$ is parameterized. ■

Assume:

- (1) There exists a finite vector α_w^* such that $P_{obs}(S_w(X; \alpha_w^*)) = 0$ where $\mathbb{P}_n S_w(X; \alpha_w)$ is the derivative with respect to α_w of (5.5) and there exists a finite vector, α_g^* such that $P_{obs}(S_g(X; \alpha_g^*)) = 0$ where $\mathbb{P}_n S_g(X; \alpha_g)$ is given in (5.6). $\mu(\beta_0, Z)$ is equal to (4.4); that is, the parameterization of (4.4) in terms of β is correct. If $\bar{\pi}_K$ is correctly parameterized then put $\alpha_w^* = \alpha_{w0}$, the true parameter in the observational distribution. Similarly if $\{g_1, \dots, g_K\}$ are correctly parameterized then $\alpha_g^* = \alpha_{g0}$, the true parameter in the observational distribution.
- (2) Let Θ be neighborhood of $(\alpha_w^*, \alpha_g^*, \beta_0)$. The class of functions

$$\left\{ \frac{\partial}{\partial \alpha_g} S_\beta(X; \alpha_w, \alpha_g, \beta), \frac{\partial}{\partial \alpha_w} S_\beta(X; \alpha_w, \alpha_g, \beta), \frac{\partial}{\partial \beta} S_\beta(X; \alpha_w, \alpha_g, \beta), \frac{\partial}{\partial \alpha_w} S_w(X; \alpha_w) \right. \\ \left. \frac{\partial}{\partial \alpha_g} S_g(X; \alpha_g), S_w(X; \alpha_w)^{\otimes 2}, S_g(X; \alpha_g)^{\otimes 2}, S_\beta(X; \gamma, \alpha_w, \alpha_g, \beta)^{\otimes 2} \text{ for } (\alpha_w, \alpha_g, \beta) \in \Theta \right\}$$

is a Glivenko-Cantelli class. (For vector V , $V^{\otimes 2} = VV^T$.)

- (3) Assume that $I_w = E_{obs} \frac{\partial}{\partial \alpha_w} S_w(X; \alpha_w)$ is invertible at $\alpha_w = \alpha_w^*$, $I_g = E_{obs} \frac{\partial}{\partial \alpha_g} S_g(X; \alpha_g)$ is invertible at $\alpha_g = \alpha_g^*$ and assume that $B = E_{obs} \dot{\mu}(\beta_0, Z) \dot{\mu}(\beta_0, Z)^T$ is invertible.

Lemma 8.3. Suppose that either $\bar{\pi}_K$ or the g_t 's are correctly parameterized. Assuming 1) through 3), there exists a sequence of $\hat{\beta}$ solutions to (5.7) for which $\sqrt{n}(\hat{\beta} - \beta_0)$ converges in distribution to a multivariate normal with mean zero and var-covariance matrix, $B^{-1}\Sigma B^{-1}$ where

$$\Sigma = E_{obs} \left(S_\beta(X; \alpha_w^*, \alpha_g^*, \beta_0) + S_{\beta, \alpha_w} I_w^{-1} S_w(X; \alpha_w^*) + S_{\beta, \alpha_g} I_g^{-1} S_g(X; \alpha_g^*) \right)^{\otimes 2}$$

where $S_{\beta, \alpha_w} = E_{obs} \frac{\partial}{\partial \alpha_w} S_{\beta}(X; \alpha_w, \alpha_g^*, \beta_0) |_{\alpha_w = \alpha_w^*}$ and

$S_{\beta, \alpha_g} = E_{obs} \frac{\partial}{\partial \alpha_g} S_{\beta}(X; \alpha_w^*, \alpha_g, \beta_0) |_{\alpha_g = \alpha_g^*}$. Furthermore $\hat{B}^{-1} \hat{\Sigma} \hat{B}^{-1}$ is a consistent estimator of $B^{-1} \Sigma B^{-1}$.

Proof. Suppose that $\alpha_w^* = \alpha_{w0}$. Then as noted in the discussion concerning (8.2) we have that

$$E_{obs} S_{\beta}(X, \alpha_{w0}, \alpha_g^*, \beta_0) = E_{obs} W_{\bar{p}_K}(\bar{A}_K, \bar{L}_{K-1}; \alpha_{w0}) \dot{\mu}(\beta_0, Z)(Y - \mu(\beta_0, Z)).$$

This is zero since $\mu(\beta_0, Z)$ is equal to (4.4). Alternately suppose that $\alpha_g^* = \alpha_{g0}$. Then as noted in the discussion concerning (8.3) we have that

$$E_{obs} S_{\beta}(X, \alpha_w^*, \alpha_{g0}, \beta_0) = E_{obs} \dot{\mu}(\beta_0, Z) \left(\sum_{a_1} g_1(a_1, L_0; \alpha_{g0}) p_1(a_1 | S_0) - \mu(\beta_0, Z) \right).$$

As before this is zero since $\mu(\beta_0, Z)$ is equal to (4.4) (recall that (4.4) can be written as in (5.8)).

Proofs that there exists a consistent sequence of solutions, $\hat{\alpha}_w \xrightarrow{P} \alpha_w^*$ to $\mathbb{P}_n(S_w(X; \alpha_w) = 0$ and a consistent sequence of solutions, $\hat{\alpha}_g \xrightarrow{P} \alpha_g^*$ to $\mathbb{P}_n(S_g(X; \alpha_g) = 0$ are very similar and use assumptions 1), 2) and 3) along with lemma 2 of Aitchison and Silvey (1958). Next using existence of consistent $\hat{\alpha}_w$ and $\hat{\alpha}_g$ and again using Aitchison and Silvey's lemma 2 along with assumptions 1), 2), 3) and 5) one can show there exist a consistence solution, $\hat{\beta} \xrightarrow{P} \beta_0$. These proofs are omitted.

To prove asymptotic normality use Taylor series arguments, along with the Glivenko-Cantelli property in 2) and the invertibility in 3), to deduce that

$$\sqrt{n}(\hat{\alpha}_w - \alpha_w^*) = I_w^{-1} \sqrt{n} \mathbb{P}_n(S_w(X; \alpha_w^*)) + o_P(1)$$

and that

$$\sqrt{n}(\hat{\alpha}_g - \alpha_g^*) = I_g^{-1} \sqrt{n} \mathbb{P}_n(S_g(X; \alpha_g^*)) + o_P(1).$$

Next a Taylor series yields,

$$\begin{aligned} 0 &= \mathbb{P}_n S_{\beta}(X, \hat{\alpha}_w, \hat{\alpha}_g, \hat{\beta}) \\ &= \mathbb{P}_n S_{\beta}(X, \alpha_w^*, \alpha_g^*, \beta_0) \\ &\quad + \mathbb{P}_n \frac{\partial}{\partial \alpha_w} S_{\beta}(X; \alpha_w, \hat{\alpha}_g, \hat{\beta}) |_{\alpha_w = \alpha_w^{**}} (\hat{\alpha}_w - \alpha_w^*) \\ &\quad \text{where } \alpha_w^{**} \text{ is a vector of values intermediate between } \hat{\alpha}_w \text{ and } \alpha_w^* \\ &\quad + \mathbb{P}_n \frac{\partial}{\partial \alpha_g} S_{\beta}(X; \alpha_w, \alpha_g, \beta) |_{\alpha_g = \alpha_g^{**}} (\hat{\alpha}_g - \alpha_g^*) \\ &\quad \text{where } \alpha_g^{**} \text{ is a vector of values intermediate between } \hat{\alpha}_g \text{ and } \alpha_g^* \\ &\quad + \mathbb{P}_n \frac{\partial}{\partial \beta} S_{\beta}(X; \alpha_w^*, \alpha_g^*, \beta^{**}) (\hat{\beta} - \beta_0) \\ &\quad \text{where } \beta^{**} \text{ is a vector of values intermediate between } \hat{\beta} \text{ and } \beta_0. \end{aligned}$$

Note from the previous discussion, the first term in the above expansion has mean zero. Next apply the Glivenko-Cantelli property in 2), along with the invertibility in 3) to yield,

$$\sqrt{n}(\hat{\beta} - \beta_0) = B^{-1} \mathbb{P}_n \left(S_\beta(X; \alpha_w^*, \alpha_g^*, \beta_0) + S_{\beta, \alpha_w} I_w^{-1} S_w(X; \alpha_w^*) + S_{\beta, \alpha_g} I_g^{-1} S_g(X; \alpha_g^*) \right) + o_P(1). \blacksquare$$

To show that $\hat{B}^{-1} \hat{\Sigma} \hat{B}^{-1}$ converges in probability to $B^{-1} \Sigma B^{-1}$, we may use the asymptotic normality result along with 2) and 3).

■

Given a parametrization of \bar{p}_K in terms of the parameter, γ , define

$$S_{num}(X; \gamma) = \frac{\partial}{\partial \gamma} \log p_t(A_t | S_{t-1}; \gamma).$$

Define

$$W_{\bar{p}_t}(\bar{A}_K, \bar{L}_{K-1}; \gamma, \alpha_w) = \prod_{j=1}^t \frac{p_j(A_j | S_{j-1}; \gamma)}{\pi_j(A_j | \bar{L}_{j-1}, \bar{A}_{j-1}; \alpha_w)}$$

and

$$S_g(X; \gamma, \alpha_g) = (Y - g_K(\bar{A}_K, \bar{L}_{K-1}; \alpha_g)) \frac{\partial}{\partial \alpha_g} g_K(\bar{A}_K, \bar{L}_{K-1}; \alpha_g) +$$

$$\sum_{t=1}^{K-1} \mathbb{P}_n \left(\sum_{a_{t+1}} p_{t+1}(a_{t+1} | S_t; \gamma) g_{t+1}(a_{t+1}, \bar{A}_t, \bar{L}_t; \alpha_g) - g_t(\bar{A}_t, \bar{L}_{t-1}; \alpha_g) \right) \frac{\partial}{\partial \alpha_g} g_t(\bar{A}_t, \bar{L}_{t-1}; \alpha_g). \blacksquare$$

And define,

$$\begin{aligned} S_\beta(X; \alpha_w, \alpha_g, \gamma, \beta) &= W_{\bar{p}_K}(\bar{A}_K, \bar{L}_{K-1}; \hat{\alpha}_w) \dot{\mu}(\beta, Z) (Y - \mu(\beta, Z)) \\ &\quad - \mathbb{P}_n \sum_{t=1}^K W_{\bar{p}_t}(\bar{A}_t, \bar{L}_{t-1}; \hat{\alpha}_w) \dot{\mu}(\beta, Z) (g_t(\bar{A}_t, \bar{L}_{t-1}; \hat{\alpha}_g) - \mu(\beta, Z)) \\ &+ \mathbb{P}_n \sum_{t=1}^K \sum_{a_t} \pi_t(a_t | \bar{A}_{t-1}, \bar{L}_{t-1}; \hat{\alpha}_w) W_{\bar{p}_t}(a_t, \bar{A}_{t-1}, \bar{L}_{t-1}; \hat{\alpha}_w) \dot{\mu}(\beta, Z) \\ &\quad (g_t(a_t, \bar{A}_{t-1}, \bar{L}_{t-1}; \hat{\alpha}_g) - \mu(\beta, Z)) \end{aligned}$$

Define

$$\begin{aligned} U(X) &= S_\beta(X; \gamma^*, \alpha_w^*, \alpha_g^*, \beta_0^*) + S_{\beta, w} I_w^{-1} S_w(X; \alpha_w^*) \\ &\quad + S_{\beta, g} I_g^{-1} (S_g(X; \gamma^*, \alpha_g^*) + S_{g, num} I_{num}^{-1} S_{num}(X; \gamma^*)) \\ &\quad + S_{\beta, num} I_{num}^{-1} S_{num}(X; \gamma^*), \end{aligned}$$

where $S_{\beta, w} = E_{obs} \frac{\partial}{\partial \alpha_w} S_\beta(X; \gamma, \alpha_w, \alpha_g^*, \beta_0^*) |_{\alpha_w = \alpha_w^*}$, $S_{\beta, g} = E_{obs} \frac{\partial}{\partial \alpha_g} S_\beta(X; \gamma, \alpha_w^*, \alpha_g, \beta_0^*) |_{\alpha_g = \alpha_g^*}$, and $S_{g, num} = E_{obs} \frac{\partial}{\partial \gamma} S_g(X; \gamma, \alpha_g^*) |_{\gamma = \gamma^*}$. Similarly let $\hat{\Sigma}^* = \mathbb{P}_n \hat{U}(X) \otimes^2$ where

$$\begin{aligned} \hat{U}(X) &= S_\beta(X; \hat{\gamma}, \hat{\alpha}_w, \hat{\alpha}_g, \hat{\beta}) + \hat{S}_{\beta, w} \hat{I}_w^{-1} S_w(X; \hat{\alpha}_w) \\ &\quad + \hat{S}_{\beta, g} \hat{I}_g^{-1} (S_g(X; \hat{\gamma}, \hat{\alpha}_g) + \hat{S}_{g, num} \hat{I}_{num}^{-1} S_{num}(X; \hat{\gamma}^*)) \\ &\quad + \hat{S}_{\beta, num} \hat{I}_{num}^{-1} S_{num}(X; \hat{\gamma}^*) \end{aligned}$$

and $\hat{S}_{\beta,w} = \mathbb{P}_n \frac{\partial}{\partial \alpha_w} S_\beta(X; \hat{\gamma}, \alpha_w, \hat{\alpha}_g, \hat{\beta})|_{\alpha_w = \hat{\alpha}_w}$, $\hat{S}_{\beta,g} = \mathbb{P}_n \frac{\partial}{\partial \alpha_g} S_\beta(X; \hat{\gamma}, \hat{\alpha}_w, \alpha_g, \hat{\beta})|_{\alpha_g = \hat{\alpha}_g}$, and $\hat{S}_{g,num} = \mathbb{P}_n \frac{\partial}{\partial \gamma} S_g(X; \gamma, \hat{\alpha}_g)|_{\gamma = \hat{\gamma}^*}$.

(1') There exists a finite vector α_w^* such that $P_{obs}(S_w(X; \alpha_w^*)) = 0$ and there exists a finite vector, α_g^* such that $P_{obs}(S_g(X; \alpha_g^*)) = 0$. There exists a finite vector γ^* such that $P_{obs}(S_{num}(X; \gamma^*)) = 0$ and also a finite vector β_0^* for which $\mu(\beta_0^*, Z) =$

$$\sum_{\bar{a}_K \in \mathcal{A}} \int_{\bar{l}_{K-1}} \int_y y f_Y(y | \bar{a}_K, \bar{v}_{K-1}, \bar{s}_{K-1}) \prod_{t=1}^K p_t(a_t | s_{t-1}; \gamma^*) \prod_{t=1}^{K-1} f_t(l_t | \bar{a}_t, \bar{l}_{t-1}) f_0(l_0 | z) dy d\bar{l}_{K-1}.$$

Suppose there exists a finite vector, α_g^* such that $P_{obs}(S_g(X; \gamma^*, \alpha_g^*)) = 0$. If $\bar{\pi}_K$ is correctly parameterized then put $\alpha_w^* = \alpha_{w0}$, the true parameter in the observational distribution. Similarly if $\{g_1, \dots, g_K\}$ are correctly parameterized then $\alpha_g^* = \alpha_{g0}$, the true parameter in the observational distribution.

The parameter α_w is estimated as before. Calculate $\hat{\gamma}$ by solving $0 = \mathbb{P}_n S_{num}(X; \gamma)$. Calculate $\hat{\alpha}_g$ by solving $0 = \mathbb{P}_n S_g(X; \hat{\gamma}, \alpha_g)$. Lastly estimate β^* by solving $0 = \mathbb{P}_n S_\beta(X; \hat{\alpha}_w, \hat{\alpha}_g, \hat{\gamma}, \beta)$ for $\hat{\beta}$.

(2') Let Θ be neighborhood of $(\gamma^*, \alpha_w^*, \alpha_g^*, \beta_0^*)$. The class of functions

$$\left\{ \begin{aligned} & \frac{\partial}{\partial \alpha_g} S_\beta(X; \gamma, \alpha_w, \alpha_g, \beta), \quad \frac{\partial}{\partial \alpha_w} S_\beta(X; \gamma, \alpha_w, \alpha_g, \beta), \quad \frac{\partial}{\partial \beta} S_\beta(X; \gamma, \alpha_w, \alpha_g, \beta), \\ & \frac{\partial}{\partial \gamma} S_\beta(X; \gamma, \alpha_w, \alpha_g, \beta), \quad \frac{\partial}{\partial \alpha_g} S_g(X; \gamma, \alpha_g), \quad \frac{\partial}{\partial \gamma} S_g(X; \gamma, \alpha_g), \quad \frac{\partial}{\partial \gamma} S_{num}(X; \gamma), \\ & \frac{\partial}{\partial \alpha_w} S_w(X; \alpha_w), \quad S_w(X; \alpha_w)^{\otimes 2}, \quad S_g(X; \gamma, \alpha_g)^{\otimes 2}, \quad S_\beta(X; \gamma, \alpha_w, \alpha_g, \beta_0)^{\otimes 2}, \\ & S_{num}(X; \gamma)^{\otimes 2} \quad \text{for } (\gamma, \alpha_w, \alpha_g, \beta) \in \Theta \end{aligned} \right\}$$

is a Glivenko-Cantelli class. (For vector V , $V^{\otimes 2} = VV^T$.)

(3') Assume that $I_w = E_{obs} \frac{\partial}{\partial \alpha_w} S_w(X; \alpha_w)$ is invertible at $\alpha_w = \alpha_w^*$, $I_g = E_{obs} \frac{\partial}{\partial \alpha_g} S_g(X; \alpha_g)$ is invertible at $\alpha_g = \alpha_g^*$, $I_{num} = E_{obs} \frac{\partial}{\partial \gamma} S_{num}(X; \gamma)$ is invertible at $\gamma = \gamma^*$ and assume that $B = E_{obs} \dot{\mu}(\beta_0^*, Z) \dot{\mu}(\beta_0^*, Z)^T$ is invertible.

Lemma 8.4. Suppose that either $\bar{\pi}_K$ or the g_t 's are correctly parameterized. Assuming 1') through 3'), there exists a sequence of $\hat{\beta}$ solutions to (5.7) for which $\sqrt{n}(\hat{\beta} - \beta_0^*)$ converges in distribution to a multivariate normal with mean zero and var-covariance matrix, $B^{-1} \Sigma^* B^{-1}$ where $\Sigma^* = E_{obs} (U(X))^{\otimes 2}$.

Furthermore $\hat{B}^{-1} \hat{\Sigma}^* \hat{B}^{-1}$ is a consistent estimator of $B^{-1} \Sigma^* B^{-1}$.

Proof. The proof is similar to that of the previous lemma and is omitted. ■

9. REFERENCES

- [1] Aitchison, J. and Silvey, S.D., (1958). Maximum-Likelihood Estimation of Parameters Subject to Restraints. *Ann. Math. Statist.* **29**, 813–828.
- [2] Angrist, J.D. Imbens, G.W. and Rubin, D.B., (1996). Identification of causal effects using instrumental variables. *Journal of the American Statistical Association* **91**, 434–471.
- [3] Bickel, P., C. Klaassen, Y. Ritov and Wellner, J., (1993). *Efficient and adaptive estimation for semiparametric models*. Johns Hopkins University Press, Baltimore.
- [4] Bierman, K.L., M.T. Greenberg and The Conduct Problems Prevention Research Group, (1996) Social skill training in the FAST Track program. *Preventing childhood disorders, substance abuse, and delinquency*, 65–89, (eds: R. DeV. Peters, R.J. McMahon). Sage, Newbury Park, CA.
- [5] Borhani, N.O., W.B. Applegate, J.A. Cutler, B.R. Davis, C.D. Furberg, E. Lakatos, L. Page, H.M. Perry, W.M. Smith and Probstfield, J.L., (1991) Part 1: Rationale and Design. *Hypertension* **17(suppl II)**, II-2–II-15.
- [6] Conduct Problems Prevention Research Group, (1992). A developmental and clinical model for the prevention of conduct disorders: The FAST Track Program. *Development and Psychopathology* **4**, 509–528.
- [7] Conduct Problems Prevention Research Group, (1999a). Initial impact of the Fast Track prevention trial for conduct problems: I. The high-risk sample. *to appear in the Journal of Consulting and Clinical Psychology*.
- [8] Conduct Problems Prevention Research Group, (1999b). Initial impact of the Fast Track prevention trial for conduct problems: II. Classroom effects *to appear in the Journal of Consulting and Clinical Psychology*.
- [9] Cooperative Research Group, (1988). Rationale and design of a randomized clinical on prevention of stroke in isolated systolic hypertension. *Journal of Clinical Epidemiology* **41**, 1197–1208.
- [10] Cox, D.R. (1958). *The Design and Planning of Experiments*. Chapman and Hall, London.
- [11] Hernán, M.A., B. Brumback, and Robins, J.M., (2000). Marginal structural models to estimate the causal effect of zidovudine on the survival of HIV-positive men. *Epidemiology* **11**, 561-570.
- [12] Holland, P., (1986). Statistics and Causal Inference. *Journal of the American Statistical Association* **81**, 945–961.
- [13] Manski, C.F., (1995). *Identification Problems in the Social Sciences*. Harvard University Press, Cambridge, MA.
- [14] McMahon, R.J., N. Slough and Conduct Problems Prevention Research Group, (1996). Family-based intervention in the FAST Track Program. *Preventing childhood disorders, substance abuse, and delinquency*, 65–89, (eds: R. DeV. Peters, R.J. McMahon).

Sage, Newbury Park, CA.

- [15] Neyman, J. (1923). On the application of probability theory to agricultural experiments. Translated in *Statistical Science*, bf 5, 465–480 (1990).
- [16] Robins, J.M. (1986). A new approach to causal inference in mortality studies with sustained exposure periods—application to control of the healthy worker survivor effect. *Computers and Mathematics with Applications* **14**, 1393–1512.
- [17] Robins, J.M. (1987). Addendum to "A new approach to causal inference in mortality studies with sustained exposure periods - Application to control of the healthy worker survivor effect." *Computers and Mathematics with Applications* **14**, 923–945.
- [18] Robins, J.M. (1989). The Analysis of Randomized and Nonrandomized AIDS Treatment Trials Using a New Approach to Causal Inference in Longitudinal Studies. *Health Service Research Methodology: A Focus on AIDS*, (eds: L. Sechrest, H. Freeman, A. Mulley). NCHSR, U.S. Public Health Service, 113–159.
- [19] Robins, J.M. (1993). Information recovery and bias adjustment in proportional hazards regression analysis of randomized trials using surrogate markers. *Proceedings of the Biopharmaceutical Section, American Statistical Association*, 24–33.
- [20] Robins, J.M. (1997). Causal Inference from Complex Longitudinal Data. *Latent Variable Modeling and Applications to Causality. Lecture Notes in Statistics (120)*, 69–117, (eds: M. Berkane). Springer-Verlag, Inc, New York.
- [21] Robins, J.M. (1998). Marginal structural models. *1997 Proceedings of the American Statistical Association. Section on Bayesian Statistical Science*, 1–10.
- [22] Robins, J.M. (1999). Testing and estimation of direct effects by reparameterizing directed acyclic graphs with structural nested models. *Computation, Causation, and Discovery.*, (eds: C. Glymour and G. Cooper). AAAI Press/The MIT Press, Menlo Park, CA & Cambridge, MA, 349-405.
- [23] Robins, J.M. (2000). Robust estimation in sequentially ignorable missing data and causal inference models. *to appear, Proceedings of the American Statistical Association*
- [24] Robins, J.M., S. Greenland, and Hu, F.C.,(1999b) Estimation of the Causal Effect of a Time-Varying Exposure on the Marginal Mean of a Repeated Binary Outcome. *Journal of the American Statistical Association* **94**, 687–700.
- [25] Robins, J.M., M.A. Hernán, and Brumback, B. (2000). Marginal structural models and causal inference in epidemiology. *Epidemiology* **11**, 550–560.
- [26] Robins, J.M. and Rotnitzky A. (1992). Recovery of information and adjustment for dependent censoring using surrogate markers. *AIDS Epidemiology - Methodological Issues*. (eds: Jewell N., Dietz K., Farewell V.) Birkhäuser, Boston, MA: 297–331.
- [27] Robins, J.M. and Rotnitzky A. (1995). Semiparametric efficiency in multivariate regression models with missing data. *Journal of the American Statistical Association* **90**, 122–129.

- [28] Robins, J.M., A. Rotnitzky and Scharfstein, D.O. (1999a). Sensitivity Analysis for Selection Bias and Unmeasured Confounding in Missing Data and Causal Inference Models. *Statistical Methods in Epidemiology*, 1–92, (eds: E. Halloran). Springer-Verlag, Inc , New York.
- [29] Robins, J.M. and Wasserman, L. (1997). Estimation of Effects of Sequential Treatments by Reparameterizing Directed Acyclic Graphs. *Proceedings of the Thirteenth Conference on Uncertainty in Artificial Intelligence*, 409–42-, (eds: D. Geiger, P. Shenoy). Morgan Kaufmann, San Francisco.
- [30] Rubin, D.B., (1978). Bayesian Inference for Causal Effects: The Role of Randomization. *The Annals of Statistics*, **6**, 34–58.
- [31] Rubin, D.B., (1986). Comment: Which Ifs Have Causal Answers *Journal of the American Statistical Association* **81**, 961–962.
- [32] Scharfstein, D.O., A. Rotnitzky and Robins, J.M. (1999). Adjusting for Nonignorable Drop-Out Using Semiparametric Nonresponse Models. *Journal of the American Statistical Association* **94**, 1096–1120.